



ދިވެހިރާއްޖޭގެ ރައްކާތެރިކަމުގެ ސަބަބަށް ފުރިހަމަކުރެއްވި ދަޢުވާތަކާ ގުޅިގެން

PROTECTED BIRDS OF THE MALDIVES

2026

ދިވެހިރާއްޖޭގެ ސަރުކާރުގެ ފަރާތުން
ދަތުރުތަކާ ގުޅިގެން ދަޢުވާތަކާ ގުޅިގެން



Ministry of Climate Change,
Environment and Energy
Republic of Maldives







ޖަނަބީ ސަރުކާރުގެ ގެޒެޓް

ޖުމްހޫރީ ސަރުކާރުގެ ގެޒެޓް، ހިސާބު 2026، ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި

ހިސާބު 2026،
ޖަނަބީ ސަރުކާރުގެ ގެޒެޓް،
ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި

Ministry of Climate Change, Environment and Energy (2026),
Protected Birds of the Maldives, Male: Ministry of Climate Change,
Environment and Energy.

978-99915-78-47-7

ޖުމްހޫރީ ސަރުކާރުގެ ގެޒެޓް، ހިސާބު 2026، ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
ހިސާބު 2026، ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި

ހިސާބު 2026،
ޖަނަބީ ސަރުކާރުގެ ގެޒެޓް،
ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި

ޖުމްހޫރީ ސަރުކާރުގެ ގެޒެޓް، ހިސާބު 2026، ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި

1	ޖުމްހޫރީ ސަރުކާރުގެ ގެޒެޓް	9	ޖުމްހޫރީ ސަރުކާރުގެ ގެޒެޓް
2	ހިސާބު 2026، ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި	10	ހިސާބު 2026، ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
3	ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި	11	ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
4	ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި	12	ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
5	ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި	13	ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
6	ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި	14	ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
7	ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި	15	ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
8	ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި		

ޖުމްހޫރީ ސަރުކާރުގެ ގެޒެޓް، ހިސާބު 2026، ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި

1	ހިސާބު 2026، ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި
2	ހިސާބު 2026، ސަފްތަވުނު ބަޔާން 15 ވަނަ ބައި

Copyright/credits

Published by:	Ministry of Climate Change, Environment and Energy
Copyright:	Government of Maldives
Citation:	Ministry of Climate Change, Environment and Energy (2026), Protected Birds of the Maldives, Male': Ministry of Climate Change, Environment and Energy.
ISBN:	978-99915-78-47-7
Author:	Ministry of Climate Change, Environment and Energy
Illustrations:	Open Sky Group Private Limited and Ismail Murushid
Dhivehi translations:	Kelavehi - Ifham Hassan Zareer and Hassan Nidham
Cover Page:	Open Sky Group Private Limited
Project Lead:	Ministry of Climate Change, Environment and Energy - Mohamed Zahir (Meemu Zaviyani) Director General

Contributors (Staff of Ministry of Climate Change, Environment and Energy):

- | | |
|--------------------------|--------------------------|
| 1. Fathimath Mohamed | 9. Fawaz Anwar |
| 2. Umaira Ahmed | 10. Fathimath Usra |
| 3. Mariyam Fazna | 11. Nafha Abbas |
| 4. Ali Nizar | 12. Rayya Hussain |
| 5. Ilham Atho Mohamed | 13. Fathina Ibrahim |
| 6. Mohamed Naaif | 14. Zulkifli Shah Ismail |
| 7. Hawwa Junainath | 15. Aminath Maiha Hameed |
| 8. Khadeeja Jumana Siraj | |

Voluntary Contributors:

1. Amir Rasheed
2. Hanan Ahmed



دعوتِ اسلامی کی راہِ راستہ پر چلنا اور اللہ کے فضل سے نفع اٹھانا

دعوتِ اسلامی کی راہِ راستہ پر چلنا اور اللہ کے فضل سے نفع اٹھانا، اس کا مطلب ہے کہ ہم اللہ کے فضل سے نفع اٹھانے کے لیے اللہ کی راہِ راستہ پر چلیں اور اللہ کے فضل سے نفع اٹھانے کے لیے اللہ کی راہِ راستہ پر چلیں۔

دعوتِ اسلامی کی راہِ راستہ پر چلنا اور اللہ کے فضل سے نفع اٹھانا، اس کا مطلب ہے کہ ہم اللہ کے فضل سے نفع اٹھانے کے لیے اللہ کی راہِ راستہ پر چلیں اور اللہ کے فضل سے نفع اٹھانے کے لیے اللہ کی راہِ راستہ پر چلیں۔

دعوتِ اسلامی کی راہِ راستہ پر چلنا اور اللہ کے فضل سے نفع اٹھانا، اس کا مطلب ہے کہ ہم اللہ کے فضل سے نفع اٹھانے کے لیے اللہ کی راہِ راستہ پر چلیں اور اللہ کے فضل سے نفع اٹھانے کے لیے اللہ کی راہِ راستہ پر چلیں۔

دعوتِ اسلامی کی راہِ راستہ پر چلنا اور اللہ کے فضل سے نفع اٹھانا، اس کا مطلب ہے کہ ہم اللہ کے فضل سے نفع اٹھانے کے لیے اللہ کی راہِ راستہ پر چلیں اور اللہ کے فضل سے نفع اٹھانے کے لیے اللہ کی راہِ راستہ پر چلیں۔

دعوتِ اسلامی کی راہِ راستہ پر چلنا اور اللہ کے فضل سے نفع اٹھانا، اس کا مطلب ہے کہ ہم اللہ کے فضل سے نفع اٹھانے کے لیے اللہ کی راہِ راستہ پر چلیں اور اللہ کے فضل سے نفع اٹھانے کے لیے اللہ کی راہِ راستہ پر چلیں۔

اللہ کے فضل سے نفع اٹھانا
دعوتِ اسلامی کی راہِ راستہ پر چلنا



FOREWORD

MINISTER OF CLIMATE CHANGE, ENVIRONMENT AND ENERGY

The natural environment and biodiversity of the Maldives is our most important asset. Just as our ancestors entrusted us with our natural heritage, it is our responsibility to pass them on to future generations.

The rich natural environments of the Maldives serve as habitats for a great number of migratory birds from different parts of the world and the Maldives remains an important stopover and destination for these species; therefore, it is vital that our ecosystems remain a safe haven for them.

Migratory birds are now protected by law in the Maldives, and it is our collective responsibility to ensure a safe environment for them.

This book, “Protected Birds of the Maldives” has been compiled by the staff of the Environmental Management and Conservation Department of this Ministry, under the supervision of the department’s Director General, Mr. Mohamed Zahir. It contains illustrations of and information on both commonly spotted and migratory birds of the Maldives which are protected under the Environmental Protection and Preservation Act of the Maldives. Our hope is that the information presented in this book will help increase understanding of avifauna and inspire greater interest in their conservation.

I wish to extend my sincere appreciation to everyone who contributed to this book. May Almighty Allah bless our efforts to protect and preserve the Maldives’ natural environment and biodiversity for generations to come.

Ali Shareef

Minister



ادبیات

دعوتِ اسلامی کی روشنیوں میں زندگی - ڈاکٹر محمد رفیع (پروفیسر)

"دعوتِ اسلامی کی روشنیوں میں زندگی" کے نام سے ایک کتاب لکھی گئی ہے جس میں دعوتِ اسلامی کی روشنیوں میں زندگی کے مختلف پہلوؤں پر روشنی ڈالی گئی ہے۔ اس کتاب میں دعوتِ اسلامی کی روشنیوں میں زندگی کے مختلف پہلوؤں پر روشنی ڈالی گئی ہے۔ اس کتاب میں دعوتِ اسلامی کی روشنیوں میں زندگی کے مختلف پہلوؤں پر روشنی ڈالی گئی ہے۔

اس کتاب میں دعوتِ اسلامی کی روشنیوں میں زندگی کے مختلف پہلوؤں پر روشنی ڈالی گئی ہے۔ اس کتاب میں دعوتِ اسلامی کی روشنیوں میں زندگی کے مختلف پہلوؤں پر روشنی ڈالی گئی ہے۔ اس کتاب میں دعوتِ اسلامی کی روشنیوں میں زندگی کے مختلف پہلوؤں پر روشنی ڈالی گئی ہے۔

اس کتاب میں دعوتِ اسلامی کی روشنیوں میں زندگی کے مختلف پہلوؤں پر روشنی ڈالی گئی ہے۔ اس کتاب میں دعوتِ اسلامی کی روشنیوں میں زندگی کے مختلف پہلوؤں پر روشنی ڈالی گئی ہے۔ اس کتاب میں دعوتِ اسلامی کی روشنیوں میں زندگی کے مختلف پہلوؤں پر روشنی ڈالی گئی ہے۔

ڈاکٹر محمد رفیع (پروفیسر)
دعوتِ اسلامی کی روشنیوں میں زندگی



FOREWORD

DIRECTOR GENERAL- MOHAMED ZAHIR

The main objective of this book ‘Protected Birds of the Maldives’ is to increase awareness of and appreciation for bird species that we see across the Maldives, as well as the natural environments that these birds reside in. These birds not only enhance the beauty of our ecosystems; the safety that our ecosystems provide for migratory birds is also a testament to the interconnectedness and reliance between the ecosystems of the Maldives and global biodiversity.

Most of the bird species seen in the Maldives are seasonal migratory visitors that fly vast distances from different parts of the world in search of suitable habitats. The Maldives archipelago lies along important migratory routes for birds. It is therefore crucial that we learn about the birds that we see, not only because of the benefits it brings to the tourism and fisheries sectors of the Maldivian economy, but also to deepen our understanding of how changes in bird populations are linked to climate change, water and air quality, as well as the overall health of interconnected natural ecosystems.

I would like to take this opportunity to extend my heartfelt gratitude to Ms. Umaira Ahmed for her hard work in compiling and developing this book under my supervision. I wish to also thank the team who assisted Umaira in this endeavour, as well as Ms. Mariyam Fazna, Project Manager for ENDhERI, for her diligence in completing this project. I am especially grateful to Ms. Ilham Atho Mohamed, who enthusiastically supported the idea of making this book and helped turn the idea into reality.

Mohamed Zahir

Director General



සමස්ත ලෝක සුවසෑමයේ රැකවල්

"ලෝක සුවසෑමයේ රැකවල් සලකුණු" ලෙස "ජෛව විවිධත්ව සංරක්ෂණයේ ලෝක පවුල" (විද්‍යාත්මක පවුල) 2021/R-25 (ලෝක සුවසෑමයේ රැකවල් සලකුණු) මගින් සමස්ත ලෝක සුවසෑමයේ රැකවල් සලකුණු පිළිබඳව අනුමැතිය ලබා දෙන ලදී. මෙම සලකුණු මගින් ලෝක සුවසෑමයේ රැකවල් සලකුණු පිළිබඳව අනුමැතිය ලබා දෙන ලදී. මෙම සලකුණු මගින් ලෝක සුවසෑමයේ රැකවල් සලකුණු පිළිබඳව අනුමැතිය ලබා දෙන ලදී.

ජෛව විවිධත්ව සංරක්ෂණයේ ලෝක පවුල මගින් ලෝක සුවසෑමයේ රැකවල් සලකුණු පිළිබඳව අනුමැතිය ලබා දෙන ලදී. මෙම සලකුණු මගින් ලෝක සුවසෑමයේ රැකවල් සලකුණු පිළිබඳව අනුමැතිය ලබා දෙන ලදී. මෙම සලකුණු මගින් ලෝක සුවසෑමයේ රැකවල් සලකුණු පිළිබඳව අනුමැතිය ලබා දෙන ලදී.

ජෛව විවිධත්ව සංරක්ෂණයේ ලෝක පවුල මගින් ලෝක සුවසෑමයේ රැකවල් සලකුණු පිළිබඳව අනුමැතිය ලබා දෙන ලදී. මෙම සලකුණු මගින් ලෝක සුවසෑමයේ රැකවල් සලකුණු පිළිබඳව අනුමැතිය ලබා දෙන ලදී. මෙම සලකුණු මගින් ලෝක සුවසෑමයේ රැකවල් සලකුණු පිළිබඳව අනුමැතිය ලබා දෙන ලදී.

ජෛව විවිධත්ව
සංරක්ෂණයේ ලෝක පවුල



PREFACE AND ACKNOWLEDGEMENTS

This book contains information on bird species protected under the Protected Species Regulation (2025/R-25), established under the Environmental Protection and Preservation Act (4/93) of the Maldives. These species include birds that are commonly spotted in the Maldives, such as the Grey Heron, and also include migratory visitors. Hence, some of the birds included in this book may be more familiar to readers than others. Through this publication, we hope to familiarize readers with the lesser-known species and inspire interest in bird watching and the conservation of avifauna.

This book has been completed with the support and help of many; I would like to extend my sincere gratitude to everyone who helped in the development and compilation of this book. I am grateful to Amir Rasheed for the assistance he provided to the Ministry during this process. His assistance, especially in finding the Dhivehi names used for birds in different regions of the Maldives, as well as his advice on naming birds that previously lacked recorded names helped enhance the content of this book.

I would also like to thank the staff of the Environmental Management and Conservation Department who contributed to this book, including Hawwa Junainath, Khadeeja Jumana Siraj, Fawaz Anwar, Fathimath Usra, Nafha Abbas, Rayya Hussain, Fathina Ibrahim, Zulkifli Shah Ismail and Aminath Maiha Hameed. I also extend my gratitude to Hanan Ahmed for her contributions. Additional thanks go to Aishath Amal, Ahmed Raidh, Fathimath Shamra, Fathimath Nashwa, Afa Hussain, Ahmed Wisam, Thibyan Ibrahim and Ismail Ajmal, for their assistance.

I am profoundly thankful to Fathimath Mohamed for her advice and assistance during the various stages of developing this book. I also wish to acknowledge Ilham Atho Mohamed who took a special interest in this project from its inception. Finally, my deepest appreciation goes to Mohamed Zahir, Director General of the Environmental Management and Conservation department, for his leadership, guidance and support in bringing this book to fruition.

Umaira Ahmed

Content Coordinator



Contents page

ދިވެހިސަރުކާރުގެ ގެޒެޓް

Copyright/credits	03 - 02
Foreword: Ministers of Climate Change, Environment and Energy	05 - 04
Director Generals' foreword	07 - 06
Preface and Acknowledgement	09 - 08
Contents page	10 - 10
Anatomy illustrations	12 - 12
Map of Maldives	13 - 13
Introduction	17 - 14
Contents (illustrations and descriptions)	419 - 18

މާއްދާގެ ނަންބަރު 02 ގެ 2019 ވަނަ އަހަރުގެ ފެބްރުއަރީ 15 ވަނަ ދުވަހުގެ ބަންދު ދުވަހުގައި ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި
މި ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި 2019 ވަނަ އަހަރުގެ ފެބްރުއަރީ 15 ވަނަ ދުވަހުގެ ބަންދު ދުވަހުގައި ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި
މި ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި 2019 ވަނަ އަހަރުގެ ފެބްރުއަރީ 15 ވަނަ ދުވަހުގެ ބަންދު ދުވަހުގައި ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި
މި ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި 2019 ވަނަ އަހަރުގެ ފެބްރުއަރީ 15 ވަނަ ދުވަހުގެ ބަންދު ދުވަހުގައި ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި
މި ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި 2019 ވަނަ އަހަރުގެ ފެބްރުއަރީ 15 ވަނަ ދުވަހުގެ ބަންދު ދުވަހުގައި ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި
މި ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި 2019 ވަނަ އަހަރުގެ ފެބްރުއަރީ 15 ވަނަ ދުވަހުގެ ބަންދު ދުވަހުގައި ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި
މި ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި 2019 ވަނަ އަހަރުގެ ފެބްރުއަރީ 15 ވަނަ ދުވަހުގެ ބަންދު ދުވަހުގައި ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި
މި ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި 2019 ވަނަ އަހަރުގެ ފެބްރުއަރީ 15 ވަނަ ދުވަހުގެ ބަންދު ދުވަހުގައި ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ގޮތުގައި

ކަނޑުބަނޑުގެ ބަނޑުގެ ނަންމުތައް



Credit: Names taken from “Dhivehiraajjeygai... Kandumathee Dhooni” by Ahmed Shafeeq

Maldives Territory Map





Protection of Bird Species in the Maldives

Maldivians have always had a keen interest in birds. This is evidenced by fishermen using the path of seabirds to locate schools of fish, as well as different birds species being heavily featured in Maldivian folklore and children's stories passed down through generations. Yet, this fascination also led to practices negatively impacting wild populations such as the once common practice of capturing wild birds to be kept as domestic pets. Recognizing the need to safeguard bird species found in the Maldives, several major steps have been taken for the conservation of birds through environmental law.

In 1993, with the ratification of Law No: 4/93 (Environment Protection and Preservation Act of the Maldives), the foundation for the protection of species was established. On 5th June 1996, the White tern (*Gygis alba*) was declared as a protected species under this Act, making this the first bird species protected under Maldivian law. This also marked the first time a species was protected under the Environmental Protection and Preservation Act of the Maldives. The White tern which was previously known to only occur in Addu Atoll, was protected following the efforts of Mr. Ahmedfulhu (Haivakaru Ahmedfulhu), then Atoll Chief of Addu Atoll who drew attention to the decline of the species even within Addu Atoll.

Following this milestone, an additional 22 bird species were protected on 15th July 1999, a further 47 species on 22nd May 2003 and an additional 33 species in 2013.

The next significant advancement in bird conservation in the Maldives came with the introduction of Regulation No: 2014/R-169 (Regulation on Migratory Species). Under this Regulation, the capture, domestication and trade in birds and bird parts was prohibited. It also forbade the taking of eggs and any act causing harm to migratory birds or their nests. This Regulation marked a significant step in protecting migratory birds. The most recent development in bird conservation was in 2021 with the establishment of Regulation No: 2021/R-25 (Regulation on Protected Species). On the day this Regulation came into effect, a total of 207 bird species were granted legal protection. Guidelines on interactions with protected birds have also been developed under this regulation to ensure their continued conservation and welfare.

Today, these laws and regulations form the backbone of national efforts to ensure that the Maldivian environment remains a haven for bird species. This book sheds light on the beauty and diversity of avifauna found in the Maldives and serves as a reference to enhance understanding of avifauna in the Maldives.



Illustrations and Descriptions Explanatory Note

Information on the birds included in this book is derived from multiple sources, including the “Birds of the World” website, as well as Mr. Ahmed Shafeeq’s book, “Dhivehiraajjeygai... Kandumathee Dhooni”, published in 1993. Additional information has been drawn from the “Checklist of Birds of the Maldives” developed under Project REGENERATE, a Government of Maldives project implemented by the International Union for the Conservation of Nature (IUCN) with funding from USAID.

References used to identify Dhivehi names of bird species include the article “Birds of the Maldivian Islands” by J.S. Ash and Ali Shafeeq, as well as the digital book “Dhivehiraajjeyge Dhoonithah” published by Dhagatha Publishers.

Information on the abundance of bird species included in this book was obtained from the “Checklist of Birds of the Maldives” which used the following classification criteria:

Vagrant:	Just one or two records
Rare:	No more than 10 records
Uncommon:	More than 10 records but unlikely to be seen during any one week visit
Regular:	Might be seen during a one week visit (in the right atoll and season)
Common:	Should be seen during a one week visit (in the right atoll and season)

Information on the IUCN Red List Category for each species was sourced from the IUCN Red List of Threatened Species website, with the year of assessment indicated in brackets. Information on whether each species is listed under the Convention on International Trade in Endangered Species of Wild Fauna and Flora has been taken from the Checklist of CITES Species website. This information is accurate at the time of publication, however, this information may change.

During the development of this book, Dhivehi names were assigned to certain bird species for which no local names could be found in existing literature or through consultation. To formalize these names, suggestions were submitted to the Dhivehi Bas Committee. The suggested names were based on the bird’s appearance, common English name, and scientific name. The Dhivehi names presented in this book are those approved by the Dhivehi Bas Committee. It is acknowledged that some bird species, including those listed here, may be known by different names in various parts of the Maldives.

The illustrations presented in this book primarily depict male birds in their breeding plumage. There may be variations in the appearance, colouring or marking of individuals observed in the wild.



رَمَدَّ وَ مَنَدَّ وَ تَمَنَدَّ

White Wagtail
Motacilla alba



ہندوستان کے پرندوں کی فہرست

Indian Paradise-Flycatcher
Terpsiphone paradisi



ہندوستان کا پرندہ

Indian Golden Oriole

Oriolus kundoo



رِسْتَمَرِ بَرَجَوِيَّ

Indian Robin

Saxicoloides fulicatus



پرنده مهاجر

Indian Cuckooshrike

Coracina macei



درخت سبز

Indian Roller

Coracias benghalensis



ہندوستان کے پرندوں کی فہرست

Indian Cuckoo

Cuculus micropterus



۳۹

Lesser Frigatebird

Fregata ariel



މާލިއްޔާ ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ދިވެހިގެޒެޓް ގައި ބަޔާންކުރި

Maldivian Pond Heron

Ardeola greyii phillipsi

ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ޕްރޮޓެކްޓެޑް ބައިރުގެ ނަންބަރު 107 ގައި ބަޔާންކުރި ޕްރޮޓެކްޓެޑް ބައިރުގެ ނަންބަރު 107

Maldivian Pond Heron

Ardeola greyii phillipsi

Status: Breeding resident in south

Local Abundance: Common

IUCN Red List Status: Unknown

CITES Appendix: Not listed

Description: Is a relatively small wader with a distinctive, pale grey-brown body, white underparts, and a short, stout neck. It has yellow legs and a sharp bill, with a slight reddish tinge on the back during the breeding season.

ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ޕްރޮޓެކްޓެޑް ބައިރުގެ ނަންބަރު 107

ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ޕްރޮޓެކްޓެޑް ބައިރުގެ ނަންބަރު 107 ގައި ބަޔާންކުރި ޕްރޮޓެކްޓެޑް ބައިރުގެ ނަންބަރު 107 ގައި ބަޔާންކުރި ޕްރޮޓެކްޓެޑް ބައިރުގެ ނަންބަރު 107



سُرَّوْرُ مَکْرُورِیَّ

Pallid Harrier

Circus macrourus



رَبِّهِمْ وَتُحْمَرُهُمْ

Great White Pelican

Pelecanus onocrotalus



سُرَقْرَسِرْدَیْ قُوْمُ قُرْسِرِ

White-Tailed Tropicbird

Phaethon lepturus



بند رومپڈ سٹورم پیٹریل

Band-rumped Storm-petrel

Hydrobates castro



سُرْمَرِ سَاسِرِي سَاسِرِي

White-faced Storm-petrel

Pelagodroma marina



دروغہ سے بچنے کے لیے

Masked Booby

Sula dactylatra



لچ کے طوفان پتrel

Leach's Storm-petrel

Hydrobates leucorhous

ރުކުލަވަތަރު ސަރުޖަ ސަރުޖަ

Leach's Storm-petrel

Hydrobates leucorhous

Status:	-
Local Abundance:	Vagrant
IUCN Red List Status:	Vulnerable (2018)
CITES Appendix:	Not listed

Description: Small seabird that measures 18-22 cm in length. Appears uniformly dark blackish brown with grey upperparts. Has a dark crown, tail and flight feathers. Has a characteristic white patch on the rump and a broad diagonal wing bar which is pale in color.

ރުކުލަވަތަރު ސަރުޖަ ސަރުޖަ އަކީ 18-22 ސެންޓިމީޓަރު ދެގަނޑުގެ ސަރުޖަ ސަރުޖަ އެވެ. ހުދު ކަރުކަތުގެ ބައިތައް ވަނީ ފަދަ ވަރަށް ފުރިހަމަކުރެއްވި ގަވާއިދުގައި ރަތު ސަރުޖަ ސަރުޖަ ގޮތުގައި ދައްކަވާފައެވެ. ހުދު ކަރުކަތުގެ ބައިތައް ވަނީ ފަދަ ވަރަށް ފުރިހަމަކުރެއްވި ގަވާއިދުގައި ރަތު ސަރުޖަ ސަރުޖަ ގޮތުގައި ދައްކަވާފައެވެ. ހުދު ކަރުކަތުގެ ބައިތައް ވަނީ ފަދަ ވަރަށް ފުރިހަމަކުރެއްވި ގަވާއިދުގައި ރަތު ސަރުޖަ ސަރުޖަ ގޮތުގައި ދައްކަވާފައެވެ. ހުދު ކަރުކަތުގެ ބައިތައް ވަނީ ފަދަ ވަރަށް ފުރިހަމަކުރެއްވި ގަވާއިދުގައި ރަތު ސަރުޖަ ސަރުޖަ ގޮތުގައި ދައްކަވާފައެވެ.



سُرْمَرِیَیْ

Streaked Shearwater

Calonectris leucomelas



سُوتی شِیئر وِٹَر

Sooty Shearwater

Ardenna grisea

ދަރަނި ޖަލު

Sooty Shearwater

Ardenna grisea

Status:	Offshore migrant
Local Abundance:	Rare
IUCN Red List Status:	Near Threatened (2019)
CITES Appendix:	Not listed

Description: Medium-sized seabird, measuring 40–51 cm in length. Dark brown in color with silver flashes on the underwings. Has narrow wings and a quick, deep wingbeat. Has a distinctive “shearing” flight style, in which it dips from side to side on stiff wings with minimal wingbeats, often skimming the water. In poor lighting, it appears completely black; in good light, it is seen as dark brown with a silvery stripe on the underwings.

ބުނެވާނެ ސަފުދި

މަތީ ގެ ސަފުދި 40-51 ސެންޓިމީޓަރު ގެ ދިގު ސަފުދި ގަނޑު ހުރެންދެން ފެންނަނީ ދަރަނި ޖަލުގެ ބަނޑުގައި. ބަނޑު ދެކޮޅުގައި ސަފުދި ލަނޑު ފެނެއެވެ. ބަނޑުގެ ދަށުގައި ވިޔަފުހެތަކުގެ ފެންވަރެއް ހުރެއެވެ. ބަނޑުގެ ދަށުގައި ވިޔަފުހެތަކުގެ ފެންވަރެއް ހުރެއެވެ. ބަނޑުގެ ދަށުގައި ވިޔަފުހެތަކުގެ ފެންވަރެއް ހުރެއެވެ. ބަނޑުގެ ދަށުގައި ވިޔަފުހެތަކުގެ ފެންވަރެއް ހުރެއެވެ. ބަނޑުގެ ދަށުގައި ވިޔަފުހެތަކުގެ ފެންވަރެއް ހުރެއެވެ. ބަނޑުގެ ދަށުގައި ވިޔަފުހެތަކުގެ ފެންވަރެއް ހުރެއެވެ. ބަނޑުގެ ދަށުގައި ވިޔަފުހެތަކުގެ ފެންވަރެއް ހުރެއެވެ.



بیرد سہ

Black-headed Heron

Ardea melanocephala



میراپس فیلیپینس

Blue-tailed Bee-eater

Merops philippinus

Blue-tailed Bee-eater

Blue-tailed Bee-eater

Merops philippinus

Status: Northern Winter Visitor
Local Abundance: Rare
IUCN Red List Status: Least Concern (2024)
CITES Appendix: Not listed

Description: Small bird measuring 23-26 cm in length. Has overall green plumage. The face has a narrow blue patch with a black eye stripe, and a yellow and brown rufous throat with bright rufous underwings visible in flight. Has a blue tail and a black beak. Juveniles are similar but paler, with a tannish throat and shorter tail.

Blue-tailed Bee-eater
23-26 cm long. Overall green plumage. Blue patch on face with black eye stripe. Yellow and brown rufous throat with bright rufous underwings. Blue tail and black beak. Juveniles are paler with tannish throat and shorter tail.



پرنده‌های ایران

Blue-cheeked Bee-eater

Merops persicus

ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި
ބަޔާންކޮށްފައިވާ ފަދައެވެ.

Blue-cheeked Bee-eater

Merops persicus

- Status:** Irruptive
- Local Abundance:** Rare
- IUCN Red List Status:** Least concern (2024)
- CITES Appendix:** Not listed

Description: Slender bird measuring about 31 cm in length. Has elongated central tail feathers that add another 7 cm. Has a lime green body and the face has a thin black mask with aquamarine blue sides. Has a bright yellow-and-red throat patch and a black beak. Males and females are similar, but the female has shorter tail-streamers.

މިބައިގެ ފައިލު ދަތުރުކުރުމަށް ފަސޭހަ ބެލުމަށް ތިބެވޭގޮތަށް ޕްލީޒް ދަތުރުކުރިއަށް ފޯމުގެ ތެރޭގައި ބުނެދޭށެވެ.
 ފަންނު ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ. ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ.
 ފަންނު ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ. ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ.
 ފަންނު ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ. ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ.
 ފަންނު ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ. ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ.
 ފަންނު ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ. ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ.
 ފަންނު ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ. ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ.
 ފަންނު ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ. ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ފަދައެވެ.



مروہ و تورتہ

European Turtle Dove

Streptopelia turtur



مردم سبز و سبز مردم

Northern House Martin

Delichon urbicum



بڑا بھرتھ سریر

Great Bittern

Botaurus stellaris

މަކިރި
މަކިރި

Great Bittern

Botaurus stellaris

Status: -

Local Abundance: Unknown

IUCN Red List Status: Least Concern (2016)

CITES Appendix: Not listed

Description:

Largest bittern species measuring 69-81 cm in length. Has a black crowned head and nape. Has tawny-buff sides on the head and neck marked with black bars. Breast and belly has brown stripes. Back and wings are rusty brown with black streaks and mottling. Bill is a greenish-yellow color. Skin around the eyes are bluish green and eyes are yellow. Legs and feet are green with yellow highlights. Males are typically bigger than females. Juveniles appear similar to adults but are have paler plumage and are less bodily marked.

މަކިރި

މަކިރި 81-69 ސަންޓިމީޓަރުގެ މަތިން ދުނިޔޭގެ ބޮޑުވަންޑެވެ. ސަލާ ގަސް ގަހުގެ ގޮތުގައި ބަހުލުވާ ނުވަތަ ހުޅަނގުގެ ގޮތުގައި ބަހުލުވާ އުޅެއެވެ. މަކިރިގެ ފަތި ބަނޑުގައި ޅެނިވިލެނި ފަދަ ގަނޑުތަކެއް ހުރެއެވެ. ބަނޑު ބަނޑުގެ މަތީ ބައިގައި ފަދަ ގަނޑުތަކެއް ހުރެއެވެ. ބަނޑު ބަނޑުގެ ތިރީ ބައިގައި ފަދަ ގަނޑުތަކެއް ހުރެއެވެ. ބަނޑު ބަނޑުގެ ތިރީ ބައިގައި ފަދަ ގަނޑުތަކެއް ހުރެއެވެ. ބަނޑު ބަނޑުގެ ތިރީ ބައިގައި ފަދަ ގަނޑުތަކެއް ހުރެއެވެ. ބަނޑު ބަނޑުގެ ތިރީ ބައިގައި ފަދަ ގަނޑުތަކެއް ހުރެއެވެ. ބަނޑު ބަނޑުގެ ތިރީ ބައިގައި ފަދަ ގަނޑުތަކެއް ހުރެއެވެ. ބަނޑު ބަނޑުގެ ތިރީ ބައިގައި ފަދަ ގަނޑުތަކެއް ހުރެއެވެ.



دروہی سہ ماہی

Fulvous Whistling Duck

Dendrocygna bicolor



مغربی زرد کھنکھ

Western Yellow Wagtail

Motacilla flava

Western Yellow Wagtail

Western Yellow Wagtail

Motacilla flava

- Status: Northern winter passage migrant
- Local Abundance: Uncommon
- IUCN Red List Status: Least Concern (2024)
- CITES Appendix: Not listed

Description:

Small bird, about 16-17 cm long. Breeding males have a blue-grey forehead and neck, darker ear-coverts, and a white line running from the bill to the neck. Has olive-brown upperparts and dark wings with yellow edges and contrasting bright yellow underparts, often with a greenish hint and a subtle dark necklace. Breeding females are duller, with browner upperparts and paler yellow underparts, sometimes spotted around the neck. Non-breeding males are more subdued, with a yellowish rump, while non-breeding females are even paler with a more noticeable necklace. Juveniles resemble non-breeding females but are often greener.

Western Yellow Wagtail

Western Yellow Wagtail 16-17 cm long. Breeding males have a blue-grey forehead and neck, darker ear-coverts, and a white line running from the bill to the neck. Has olive-brown upperparts and dark wings with yellow edges and contrasting bright yellow underparts, often with a greenish hint and a subtle dark necklace. Breeding females are duller, with browner upperparts and paler yellow underparts, sometimes spotted around the neck. Non-breeding males are more subdued, with a yellowish rump, while non-breeding females are even paler with a more noticeable necklace. Juveniles resemble non-breeding females but are often greener.



موتاسیلا سیترولا

Citrine Wagtail

Motacilla citreola



پالید سکوپس اوول

Pallid Scops-Owl

Otus brucei

ކްރިސީ ޅަވާ ޅަވާ ޅަވާ

Pallid Scops-Owl

Otus brucei

Status:	-
Local Abundance:	Vagrant
IUCN Red List Status:	Least concern (2021)
CITES Appendix:	Appendix II

Description: Small bird measuring about 18-21 cm long. Has a pale, soft-looking appearance with short ear tufts and fine black streaks across the body. The face is a light buff color with a thin dark outline and small white “eyebrows” above its pale yellow eyes. The back is light greyish-brown with a faint yellowish tint and subtle streaks. The chest and belly are paler and more heavily marked with black. The wings and tail show faint striping, and the legs are fully feathered down to the toes. Juveniles are even paler, with shorter ear tufts and softer markings.

ކްރިސީ ޅަވާ 21-18 ސެންޓިމީޓަރުގެ ތެރޭގައި ވާ ޕެލިޑް ސްކޮޕްސް-އޮޕްލް ބަނޑު ފެނިގެން ގެންދަނީ ރާއްޖޭގެ ތެރޭގައި ހުންނަ ތަންތަނުގައެވެ. ޕެލިޑް ސްކޮޕްސް-އޮޕްލް ގެ ސަބަބުން ވަނީ ރާއްޖޭގެ ތެރޭގައި ބަނޑު ފެނިގެން ގެންދަނީ ރާއްޖޭގެ ތެރޭގައި ހުންނަ ތަންތަނުގައެވެ. ޕެލިޑް ސްކޮޕްސް-އޮޕްލް ގެ ސަބަބުން ވަނީ ރާއްޖޭގެ ތެރޭގައި ބަނޑު ފެނިގެން ގެންދަނީ ރާއްޖޭގެ ތެރޭގައި ހުންނަ ތަންތަނުގައެވެ. ޕެލިޑް ސްކޮޕްސް-އޮޕްލް ގެ ސަބަބުން ވަނީ ރާއްޖޭގެ ތެރޭގައި ބަނޑު ފެނިގެން ގެންދަނީ ރާއްޖޭގެ ތެރޭގައި ހުންނަ ތަންތަނުގައެވެ. ޕެލިޑް ސްކޮޕްސް-އޮޕްލް ގެ ސަބަބުން ވަނީ ރާއްޖޭގެ ތެރޭގައި ބަނޑު ފެނިގެން ގެންދަނީ ރާއްޖޭގެ ތެރޭގައި ހުންނަ ތަންތަނުގައެވެ. ޕެލިޑް ސްކޮޕްސް-އޮޕްލް ގެ ސަބަބުން ވަނީ ރާއްޖޭގެ ތެރޭގައި ބަނޑު ފެނިގެން ގެންދަނީ ރާއްޖޭގެ ތެރޭގައި ހުންނަ ތަންތަނުގައެވެ.



مَکْرَجَة سَوِيْر

Black-crowned Night Heron

Nycticorax nycticorax



درختوں کے پتوں کی طرح ہر پرندے کا اپنا رنگ ہے

Indian Pond Heron

Ardeola greyii greyii



نورجی سٹورٹ

Lesser Kestrel

Falco naumanni



مردو سرکے کھنڈر

Red-tailed Shrike

Lanius phoenicuroides



مجموعہ سائنس و ٹیکنالوجی

Red-tailed Tropicbird

Phaethon rubricauda

Red-tailed Tropicbird

Red-tailed Tropicbird

Phaethon rubricauda

Status:	Straggler
Local Abundance:	Rare
IUCN Red List Status:	Least Concern (2020)
CITES Appendix:	Not Listed

Description: Medium-sized seabird, measuring 44-47 cm in length . Has a mostly white body that shines with a satiny sheen and can sometimes have a pale pink tint, which fades as the bird ages. Has a striking black crescent shape around the front of the eye, and a black line extending behind the eye. The bill is long and bright red or orange with a black tip, slightly curved and serrated. The most distinctive feature is the long, slender tail feathers, which are filamentous and can be longer than the body. The wings are long and narrow and the feet are black at the tips, with the base of the legs being a light blue-grey color. Juvenile birds look quite different, with a white forecrown that sharply contrasts with the dark spots on the head and body.


Red-tailed Tropicbird (44-47 cm) is a medium-sized seabird with a mostly white body that shines with a satiny sheen and can sometimes have a pale pink tint, which fades as the bird ages. It has a striking black crescent shape around the front of the eye, and a black line extending behind the eye. The bill is long and bright red or orange with a black tip, slightly curved and serrated. The most distinctive feature is the long, slender tail feathers, which are filamentous and can be longer than the body. The wings are long and narrow and the feet are black at the tips, with the base of the legs being a light blue-grey color. Juvenile birds look quite different, with a white forecrown that sharply contrasts with the dark spots on the head and body.



سمنگه کوه سمنگه کوه / سمنگه کوه سمنگه کوه

Cinnamon Bittern

Botaurus cinnamomeus



مَرَجُورِی

Ferruginous Pochard

Aythya nyroca



مردھو کوڑھ

Ruddy Turnstone

Arenaria interpres



سرخ‌پایه

Common Redshank

Tringa totanus



میرحکم پور / سائینس سٹریٹ

Red-footed Booby

Sula sula



مجموعہ سائنس و طبیعت

Red-billed Tropicbird

Phaethon aethereus



سائڈرس ٹرن
Saunders's Tern
Sternula saundersi



نماد جزایر و سواحل جنوبی ایران / جزایر و سواحل

Little Tern
Sternula albifrons



کے سرے

Sooty Tern

Onychoprion fuscatus



پرنده مهاجر / نماند از سرچشمه سبز

Little Ringed Plover

Charadrius dubius

ދިވެހިސަރުކާރުގެ ގެޒެޓްގައި ރިކޯޑް ކުރިފައިވާ ލިޔުންތަކުގެ ދަށުން
ފަދަ ލިޔުންތަކުގެ ދަށުން

Little Ringed Plover

Charadrius dubius

Status:	Northern winter visitor
Local Abundance:	Uncommon
IUCN Red List Status:	Least Concern (2016)
CITES Appendix:	Not listed

Description: Small bird measuring 14-17 cm in length. Characterized by a bright yellow eye ring, with a narrow white line behind its black frontal bar and a distinct head pattern. The lack of an obvious wingbar in flight is a key feature for identification. Females have a brownish tinge to the black areas and a slightly narrower eye ring compared to males. The non-breeding adult has a faint breastband and a brownish-black head but the overall plumage does not change much. Juveniles are similar in appearance to non-breeding adults, but the olive-brown upperparts are edged with buff-colored fringes.

ފަދަ ލިޔުންތަކުގެ ދަށުން / ފަދަ ލިޔުންތަކުގެ ދަށުން 17-14 ސެންޓިމީޓަރުގެ ދެކެ ގަނޑުގައި ބަޔާލުވާ ލިޔުންތަކުގެ ދަށުން
 ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން
 ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން
 ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން
 ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން
 ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން
 ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން
 ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން ފަދަ ލިޔުންތަކުގެ ދަށުން



چھ سرے سر

Caspian Plover
Charadrius asiaticus



قلمرو پرندگانی ایران

Curlew Sandpiper

Calidris ferruginea

ކަލިއުވް ސަންޑްޕައިޕަރު

Curlew Sandpiper

Calidris ferruginea

Status:	Northern winter visitor
Local Abundance:	Common
IUCN Red List Status:	Vulnerable (2024)
CITES Appendix:	Not listed

Description: Slender, long-legged shorebird measuring about 21 cm in length with a gently curved black bill that is noticeably long. Has a slightly long neck, giving it a more graceful appearance than other sandpipers. In breeding plumage, it is mostly a rich brick-red color with a white vent and undertail area. The crown is streaked black, the back is mottled with reddish, black, and white, and some birds show a white chin and eyebrow stripe. The chest and belly often have white scalloped markings from feather fringes. In non-breeding plumage, it is pale grey-brown above with white underparts and some fine streaking on the chest. The face features a strong white eyebrow that contrasts with darker grey-brown lores and ear patches. Juveniles have a scaly look, with dark-centered grey-brown feathers edged in white or buff, and mostly white underparts with light chest streaking and sometimes a buff wash.

ކަލިއުވް

ކަލިއުވް ސަންޑްޕައިޕަރު ނުވަތަ ކަލިއުވް ސަންޑްޕައިޕަރު 21 ސެންޓިމީޓަރު ގެ ރަށުގައި ފެންނަނީ ރާއްޖޭގެ ބައެއް ތަންތަނުގައެވެ. ނިކެޔުން ފެށިގެން ބައެއް ތަންތަނުގައި ފެންނަނީ ޕެލެޔޮ ސަންޑްޕައިޕަރުގެ ސަންޑްޕައިޕަރުގެ ގޮތުގައި ކަލިއުވް ސަންޑްޕައިޕަރު ފެންނަނީ ރާއްޖޭގެ ބައެއް ތަންތަނުގައެވެ. ސަންޑްޕައިޕަރުގެ ބައެއް ވައްތަރުތަކުގެ ތެރެއިން ޕެލެޔޮ ސަންޑްޕައިޕަރުގެ ގޮތުގައި ކަލިއުވް ސަންޑްޕައިޕަރު ފެންނަނީ ރާއްޖޭގެ ބައެއް ތަންތަނުގައެވެ. ސަންޑްޕައިޕަރުގެ ބައެއް ވައްތަރުތަކުގެ ތެރެއިން ޕެލެޔޮ ސަންޑްޕައިޕަރުގެ ގޮތުގައި ކަލިއުވް ސަންޑްޕައިޕަރު ފެންނަނީ ރާއްޖޭގެ ބައެއް ތަންތަނުގައެވެ. ސަންޑްޕައިޕަރުގެ ބައެއް ވައްތަރުތަކުގެ ތެރެއިން ޕެލެޔޮ ސަންޑްޕައިޕަރުގެ ގޮތުގައި ކަލިއުވް ސަންޑްޕައިޕަރު ފެންނަނީ ރާއްޖޭގެ ބައެއް ތަންތަނުގައެވެ. ސަންޑްޕައިޕަރުގެ ބައެއް ވައްތަރުތަކުގެ ތެރެއިން ޕެލެޔޮ ސަންޑްޕައިޕަރުގެ ގޮތުގައި ކަލިއުވް ސަންޑްޕައިޕަރު ފެންނަނީ ރާއްޖޭގެ ބައެއް ތަންތަނުގައެވެ. ސަންޑްޕައިޕަރުގެ ބައެއް ވައްތަރުތަކުގެ ތެރެއިން ޕެލެޔޮ ސަންޑްޕައިޕަރުގެ ގޮތުގައި ކަލިއުވް ސަންޑްޕައިޕަރު ފެންނަނީ ރާއްޖޭގެ ބައެއް ތަންތަނުގައެވެ. ސަންޑްޕައިޕަރުގެ ބައެއް ވައްތަރުތަކުގެ ތެރެއިން ޕެލެޔޮ ސަންޑްޕައިޕަރުގެ ގޮތުގައި ކަލިއުވް ސަންޑްޕައިޕަރު ފެންނަނީ ރާއްޖޭގެ ބައެއް ތަންތަނުގައެވެ. ސަންޑްޕައިޕަރުގެ ބައެއް ވައްތަރުތަކުގެ ތެރެއިން ޕެލެޔޮ ސަންޑްޕައިޕަރުގެ ގޮތުގައި ކަލިއުވް ސަންޑްޕައިޕަރު ފެންނަނީ ރާއްޖޭގެ ބައެއް ތަންތަނުގައެވެ. ސަންޑްޕައިޕަރުގެ ބައެއް ވައްތަރުތަކުގެ ތެރެއިން ޕެލެޔޮ ސަންޑްޕައިޕަރުގެ ގޮތުގައި ކަލިއުވް ސަންޑްޕައިޕަރު ފެންނަނީ ރާއްޖޭގެ ބައެއް ތަންތަނުގައެވެ.



کاسریہ چمنی اریترین

Terek Sandpiper

Xenus cinereus



صَبْرِيَّيْنِ

Whimbrel

Numenius phaeopus



قندجوسرہ ابرقوئو

Bar-tailed Godwit

Limosa lapponica



کالی سر کی ایبیس

Black-headed Ibis

Threskiornis melanocephalus



تھرش

Eyebrowed Thrush

Turdus obscurus



پلیوئلیس فولوا

Pacific Golden Plover

Pluvialis fulva



پرنیوں کی دنیا

Western Reef Egret

Egretta gularis



تھوڑے سے

Eurasian Curlew

Numenius arquata



پالاس گولف

Pallas's Gull

Ichthyaetus ichthyaetus



ٲٲٲٲ ٲٲٲٲ

Yellow-legged Gull

Larus cachinnans



پرنسپل سائنس اور ٹیکنالوجی کے شعبے کے سربراہان

Peregrine Falcon

Falco peregrinus



۳۳۰ ۳۴

Wedge-Tailed Shearwater

Ardenna pacifica



تیر کبوتر

Greater Crested Tern

Thalasseus bergii (Lichtenstein, 1823)



بته بته

Common Buzzard

Buteo buteo



گارجینی

Garganey

Anas querquedula



گول بیلڈ ٹرن

Gull-Billed Tern

Gelochelidon nilotica



سندرزین

Sanderling

Calidris alba

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

Sanderling

Calidris alba

Status:	Northern winter visitor
Local Abundance:	Uncommon
IUCN Red List Status:	Least concern (2024)
CITES Appendix:	Not listed

Description: Small, plump bird, measuring around 18–20 cm in length. Commonly found along sandy coastal areas during the non-breeding season. Has a short, straight, black bill, with black legs and feet. Lacks a hallux (the rear toe), setting it apart from other sandpipers. Non-breeding adults have a white head, pale-grey upperparts, and white underparts, with a distinctive dark patch at the bend of the wing. In flight, a bold, wide white wing-stripe bordered by black is seen. In breeding adults, the plumage changes to reddish-brown or sandy-brown on the head, breast, and upperparts, which are coarsely mottled with darker and lighter shades, contrasting with the white underparts. Juveniles resemble nonbreeding adults but are darker, with black, mottled upperparts and a streaked, dark crown. Have a warm buff patch on the breast sides, streaked or mottled with black markings.

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު

މަލްދީވީ ރިޔާސަތުގެ ދަށުން ރައްކާތެރިކުރި ޕްރޮޓެކްޓެޑް ބައިރިތިތަކުގެ ލިސްޓު



پرنده مهاجر

Kentish Plover

Charadrius alexandrinus



کوریٹری

Temminck's Stint

Calidris temminckii



چوڑا نرہ پتو

Little Stint
Calidris minuta



ڈنلن

Dunlin

Calidris alpina



قلمرو پرندگانی پاکستان

Long-toed Stint

Calidris subminuta



گلابی پرنی

Collared Pratincole

Glareola pratincola

ލަކްރަލާ ޕްރެޓިންކޯލެ

Collared Pratincole

Glareola pratincola

Status: -

Local Abundance: Vagrant

IUCN Red List Status: Least concern (2016)

CITES Appendix: Not Listed

Description:

Medium sized bird measuring 22–25 cm in length. Has brown upperparts with an olive tint, and has a white rump. The long wingtips and deeply forked tail are black, while the throat is a striking ochre-yellow, bordered with black. The breast is brown, gradually shading to a white belly, and the underwings are a rich chestnut color. The bird also has a narrow but distinct white trailing edge on the flight feathers, a red bill with a black tip, and blackish legs. Non-breeding adults, have a less distinct black border around the throat, the lores (area around the beak) are paler, and the breast becomes mottled with grey-brown. Juveniles are mottled with black on the back and breast, and the throat is white without the black collar.

ބޮވިސިވި

ލަކްރަލާ ޕްރެޓިންކޯލެ ނުވަތަ ސޯޕްރެޖްޕްރަޓިންކޯލެ ޖެސިފާތައް 22-25 ސެންޓިމީޓަރު ގެ ކޮންމެ ފަރާތް ގައި ވެސް ދިނުމެއް ނެތި ކުޑަ ދަޢަޞްކުގެ ލަކްރަލާ ޕްރެޓިންކޯލެ އެވެ. ނުވަތަ ލަކްރަލާ ޕްރެޓިންކޯލެ ނުވަތަ ސޯޕްރެޖްޕްރަޓިންކޯލެ ޖެސިފާތައް 22-25 ސެންޓިމީޓަރު ގެ ކޮންމެ ފަރާތް ގައި ވެސް ދިނުމެއް ނެތި ކުޑަ ދަޢަޞްކުގެ ލަކްރަލާ ޕްރެޓިންކޯލެ އެވެ. ލަކްރަލާ ޕްރެޓިންކޯލެ ނުވަތަ ސޯޕްރެޖްޕްރަޓިންކޯލެ ޖެސިފާތައް 22-25 ސެންޓިމީޓަރު ގެ ކޮންމެ ފަރާތް ގައި ވެސް ދިނުމެއް ނެތި ކުޑަ ދަޢަޞްކުގެ ލަކްރަލާ ޕްރެޓިންކޯލެ އެވެ. ލަކްރަލާ ޕްރެޓިންކޯލެ ނުވަތަ ސޯޕްރެޖްޕްރަޓިންކޯލެ ޖެސިފާތައް 22-25 ސެންޓިމީޓަރު ގެ ކޮންމެ ފަރާތް ގައި ވެސް ދިނުމެއް ނެތި ކުޑަ ދަޢަޞްކުގެ ލަކްރަލާ ޕްރެޓިންކޯލެ އެވެ. ލަކްރަލާ ޕްރެޓިންކޯލެ ނުވަތަ ސޯޕްރެޖްޕްރަޓިންކޯލެ ޖެސިފާތައް 22-25 ސެންޓިމީޓަރު ގެ ކޮންމެ ފަރާތް ގައި ވެސް ދިނުމެއް ނެތި ކުޑަ ދަޢަޞްކުގެ ލަކްރަލާ ޕްރެޓިންކޯލެ އެވެ. ލަކްރަލާ ޕްރެޓިންކޯލެ ނުވަތަ ސޯޕްރެޖްޕްރަޓިންކޯލެ ޖެސިފާތައް 22-25 ސެންޓިމީޓަރު ގެ ކޮންމެ ފަރާތް ގައި ވެސް ދިނުމެއް ނެތި ކުޑަ ދަޢަޞްކުގެ ލަކްރަލާ ޕްރެޓިންކޯލެ އެވެ.



نائر ورتا ہونے

Oriental Turtle Dove

Streptopelia orientalis



معمولہ پتنگانہ

Common Teal

Anas crecca



نوددی کوچک

Lesser Noddy

Anous tenuirostris



نارنگی سرخ

Short-tailed Shearwater

Ardenna tenuirostris



سکوپ قطب

Arctic Skua

Stercorarius parasiticus

Arctic Skua

Stercorarius parasiticus

Status:	Offshore, Northern Winter
Local Abundance:	Uncommon
IUCN Red List Status:	Least concern (2018)
CITES Appendix:	Not Listed

Description: Typically measures between 41-48 cm in length, with the tail streamer of breeding adults accounting for approximately 7cm of their length. Plumage variation exists in the form light, intermediate and dark morphs. Light morph adults have brown upper parts, white underparts with yellow-ish colouring on the sides of the neck. Dark morphs are sooty brown and the intermediate morphs show characteristics between the light and dark morphs. All three morphs have a characteristic white patch on the underwing and have elongated, pointed central tail feather during breeding season.



پید ویتار

Pied Wheatear

Oenanthe pleschanka



ناتر ساروچو ساتر

Short-eared Owl

Asio flammeus



نام علمی: *Passer domesticus*

House Sparrow

Passer domesticus



سائبر

Black-naped Tern

Sterna sumatrana

ދިވެހިސަރުކާރުގެ ގެޒެޓްގައި ބަޔާންކުރި މަޢުލޫމާތު ބައްލަވާލާށެވެ.

Black-naped Tern

Sterna sumatrana

Status: Breeding Resident

Local Abundance: Common

IUCN Red List Status: Least Concern (2018)

CITES Appendix: Not listed

Description: Typically measures 33–35 cm in length. It has a white crown and forehead with a contrasting black band extending from its eyes to the back of the neck. It has pale grey upperparts and white underparts with a soft rosy tint. It has a black bill, the tip of which is sometimes yellow and it has black legs. Non-breeding adults noticeably lack the black band from the eye to the neck. Juveniles have a buff colored head with hints of grey and black and have grey underparts and a black nape.

މިއަދުގެ މަދުކަށި ދިވެހިރާއްޖޭގައި ބައްލަވާލާ ދަންނަވާ ޅެއްގެ ދެވިފައިވާ އުފަންސަ ސަރުކާރުގެ ގެޒެޓްގައި ބަޔާންކުރި މަޢުލޫމާތު ބައްލަވާލާށެވެ.



سینه سفید

White-winged Tern

Chlidonias leucopterus



بُلْوِری

Bulwer's Petrel

Bulweria bulwerii



سائے تھوڑے جھوسرے

Glossy Ibis

Plegadis falcinellus



سنگھڑ

Black Kite
Milvus migrans

ދަލުކިޔުކިޔު

Black Kite

Milvus migrans

Status: -

Local Abundance: Vagrant

IUCN Red List Status: Least Concern (2021)

CITES Appendix: Appendix II

Description: Measures 46–66 cm in length. Have reddish brown plumage and a slightly forked tail which is brown in color. Juveniles have lighter colored plumage but have a more distinct pattern than adults.

ދަލުކިޔުކިޔު

ދަލުކިޔުކިޔުގެ ނިފަލު 46-66 ސެންޓިމީޓަރުގެ ތެރޭގައި ހުންނަނީ. ނިފަލުގެ ތެރޭގައި ބަލާ ހިސާބު ވަކި ވަކި ހެއްދެވި ފެންނަނީ. ދަލުކިޔުކިޔުގެ ނިފަލު ބުނު ބުނު ހުންނަނީ. ދަލުކިޔުކިޔުގެ ނިފަލު ބުނު ބުނު ހުންނަނީ. ދަލުކިޔުކިޔުގެ ނިފަލު ބުނު ބުނު ހުންނަނީ.



سینو سٹورم پیٹرل

Swinhoe's Storm-petrel

Hydrobates monorhis



سائبرچہ سریر

Black Bittern

Ixobrychus flavicollis



تیرنگ

Watercock

Gallicrex cinerea



سائبر سائبر

Brown Booby
Sula leucogaster



سینڈویچ ٹرن

Sandwich Tern

Thalasseus sandvicensis



♀، سبز، نیلا، سیاہ

♂، نیلا، سیاہ، سبز

تاکوئی
Asian Koel
Eudynamys scolopaceus



ساقی

Cotton Teal

Nettapus coromandelianus



نورجنگ فشر

Common Kingfisher

Alcedo atthis



لٹل سٹریٹ
Little Swift
Apus affinis



نِیَءَ سَرَسِیَّر

Little Egret

Egretta garzetta



نایاب سہولت

Eurasian Hobby

Falco subbuteo

ދަރުކު ފަލުކު

Eurasian Hobby

Falco Subbuteo

Status: Northern winter visitor

Local Abundance: Rare

IUCN Red List Status: Least Concern (2021)

CITES Appendix: Appendix II

Description: Medium sized bird that measures about 28-36 cm in length. Has bright red thighs and undertail feathers. Has pale cheeks and throat. Belly may be light buff-colored or whitish. Males and females appear similar but females are slightly bigger. Juveniles are browner overall with pale edges on upper feathers. Thighs and undertail are buff colored and have a pale crown.

ދަރުކު ފަލުކު

ދަރުކު ފަލުކުގެ ލަންބަރު 28-36 ސެންޓިމީޓަރު ގެ ތެރޭގައި ހުންނެވެ. ރަތް ލަނބު ފަލުކުގެ ލަންބަރު 36-28 ސެންޓިމީޓަރު ގެ ތެރޭގައި ހުންނެވެ. ދަރުކު ފަލުކުގެ ލަންބަރު 28-36 ސެންޓިމީޓަރު ގެ ތެރޭގައި ހުންނެވެ. ރަތް ލަނބު ފަލުކުގެ ލަންބަރު 36-28 ސެންޓިމީޓަރު ގެ ތެރޭގައި ހުންނެވެ. ދަރުކު ފަލުކުގެ ލަންބަރު 28-36 ސެންޓިމީޓަރު ގެ ތެރޭގައި ހުންނެވެ. ރަތް ލަނބު ފަލުކުގެ ލަންބަރު 36-28 ސެންޓިމީޓަރު ގެ ތެރޭގައި ހުންނެވެ.



نماد چھ سہ ماہی

Lesser Sand Plover

Charadrius mongolus

ލަބަ ސަރަ ސަރަ

Lesser Sand Plover

Charadrius mongolus

Status: Northern winter visitor
Local Abundance: Uncommon
IUCN Red List Status: Endangered (2023)
CITES Appendix: Not listed

Description: Typically measuring 18-21 cm in length. Plumage is primarily grey-brown and white and has a short black bill and short grey legs. Adults in non-breeding plumage have dark brown-grey heads, nape and upperparts with large brown-grey patches on the sides of the upper chest. Forehead, eyebrow, neck and underparts are white, with dark brown on the cheeks. Males in breeding plumage have a broad chestnut breast-bank outlined by black on top, chestnut forehead and black mask on the face. Females in breeding plumage are similar, but their face mask is dark grey-brown and the colouring on their crown and upper chest are duller. Juveniles are similar to non-breeding adults.

ލަބަ ސަރަ ސަރަ 21-18 ސަންޓިމީޓަރު ދާއިރާއިން ވަނީ ވަދެއި އުފެއްދުމަށް ތައްޔާރުކޮށް ހުންނަނެވެ. ވަދެއި ހަދަނީ ތިލަ ބޮޑު ސަރަ ސަރަ ގަވާއިދުގައެވެ. ވަދެއި ހަދަނީ ތިލަ ބޮޑު ސަރަ ސަރަ ގަވާއިދުގައެވެ. ވަދެއި ހަދަނީ ތިލަ ބޮޑު ސަރަ ސަރަ ގަވާއިދުގައެވެ. ވަދެއި ހަދަނީ ތިލަ ބޮޑު ސަރަ ސަރަ ގަވާއިދުގައެވެ. ވަދެއި ހަދަނީ ތިލަ ބޮޑު ސަރަ ސަރަ ގަވާއިދުގައެވެ. ވަދެއި ހަދަނީ ތިލަ ބޮޑު ސަރަ ސަރަ ގަވާއިދުގައެވެ. ވަދެއި ހަދަނީ ތިލަ ބޮޑު ސަރަ ސަރަ ގަވާއިދުގައެވެ. ވަދެއި ހަދަނީ ތިލަ ބޮޑު ސަރަ ސަރަ ގަވާއިދުގައެވެ.



نیا نیا سر / نیا سر / نیا سر

Osprey

Pandion haliaetus



قلمرو

Western Marsh Harrier

Circus aeruginosus



گڈویٹ بلیک ٹیلڈ

Black-Tailed Godwit

Limosa limosa



پرنده مهاجر

Jack Snipe

Lymnocyptes minimus



گالیناگو گالیناگو

Common Snipe
Gallinago gallinago



سوسائبل لاپوونگ

Sociable Lapwing

Vanellus gregarius



دروغہ سب سے بڑا گنہگار ہے
Cattle Egret
Bubulcus ibis



ہریہ ڈنار

Grey Heron

Ardea cinerea

ގަލަފުހި

Grey Heron

Ardea cinerea

- Status:** Breeding Resident
- Local Abundance:** Common
- IUCN Red List Status:** Least Concern (2019)
- CITES Appendix:** Not listed

Description: Large wading bird with a long neck. Has grey upperparts and white underparts. Beak is sharp and yellow in color which turns orange in breeding season. Legs are pale greyish and long. Locally known in the Maldives as "Maakana" and is one of the most easily recognisable birds.

މާކަނާ

ގަލަފުހި ނިންނަނީ ގަލަފުހި ގެ ސަރަޙައްދުގައި. ގަލަފުހި ގެ ފެންމަލު ފެންނަނީ ގަލަފުހި ގެ ސަރަޙައްދުގައި. ގަލަފުހި ގެ ފެންމަލު ފެންނަނީ ގަލަފުހި ގެ ސަރަޙައްދުގައި. ގަލަފުހި ގެ ފެންމަލު ފެންނަނީ ގަލަފުހި ގެ ސަރަޙައްދުގައި. ގަލަފުހި ގެ ފެންމަލު ފެންނަނީ ގަލަފުހި ގެ ސަރަޙައްދުގައި. ގަލަފުހި ގެ ފެންމަލު ފެންނަނީ ގަލަފުހި ގެ ސަރަޙައްދުގައި. ގަލަފުހި ގެ ފެންމަލު ފެންނަނީ ގަލަފުހި ގެ ސަރަޙައްދުގައި.



گری واگٹیل

Grey Wagtail

Motacilla cinerea



ہریہ و نیل سرخ کھنکھریہ

Spot-billed Pelican

Pelecanus philippensis



آشی وڈ سوا لو

Ashy Woodswallow

Artamus fuscus



گریبطنی کبک

Grey-bellied Cuckoo

Cacomantis passerinus



مجموعہ سائنس و طبیعت | سائنس و طبیعت

Common Moorhen

Gallinula chloropus

Common Moorhen

Common Moorhen

Gallinula chloropus

Status:	Breeds in far south
Local Abundance:	Uncommon
IUCN Red List Status:	Least Concern (2019)
CITES Appendix:	Not listed

Description: Medium sized bird measuring 30-38 cm in length. Appears mostly black from a distance. Has a noticeable red shield on the forehead and a yellow tipped red bill. Has white lines along the sides. Tail is black with a slight green shine and has white feathers underneath. Head and neck are dark grey. Has an olive-brown tinge at the neck of the neck and rest of the body is a mix of olive brown and olive green. Legs are yellow-green and eyes are bright red. Males and females appear similar but females are slightly smaller. Juveniles appear duller. Back is brown or grey. Has lighter underparts. Bill and leg colors are more muted than in adults.

Common Moorhen

Common Moorhen (Gallinula chloropus) is a medium-sized bird, 30-38 cm long. It has a black body with a red shield on the forehead and a red bill with a yellow tip. The tail is black with a slight green shine. The head and neck are dark grey, and the rest of the body is a mix of olive brown and olive green. The legs are yellow-green, and the eyes are bright red. Males and females appear similar, but females are slightly smaller. Juveniles are duller. The back is brown or grey, and the underparts are lighter. The bill and leg colors are more muted than in adults.



پرنیوں کی دنیا

Lesser Crested Tern

Thalasseus bengalensis



نورن تھیر

Northern Wheatear


Oenanthe oenanthe



رَبْوَانَا

Grey Plover

Pluvialis squatarola



دروہی سائنس اور طبیعت کی دنیا

Northern Pintail

Anas acuta

ހަލުހު ޕިންޓައިލް

Northern Pintail

Anas acuta

Status:	Northern winter visitor
Local Abundance:	Uncommon
IUCN Red List Status:	Least Concern (2019)
CITES Appendix:	Not listed

Description: Measures 50-66 cm in length. Males and females both are slim, have long narrow necks and pointed tails. Breeding males have a chocolate-brown head, neck and throat and have white stripes on the sides of the neck. Body is grey with white and yellow-buff areas. Tail coverts are black. Wings are brownish grey. Has a metallic green and bronze sheen on the speculum. Breeding females have grey-brown head and neck. Underparts are finely spotted and flanks are brownish. Wings are grey-brown with pale tips. Juveniles are dark and brown. Wings are duller and tail is brown with buff fringes.

ބަވަހު

ހަލުހު ޕިންޓައިލް ގެ ބަދަން 50-66 ސެންޓިމީޓަރު ހުރެއެވެ. ފަލިކާ ގެ ސަބަބުން ޕިންޓައިލް ގެ ނަންބަރު ދަށަންނަން ފެންނަނީ މާލެ އާއި ރާއްޖޭގެ ބަނިތަކުންނެވެ. ފަލިކާ ގެ ސަބަބުން ޕިންޓައިލް ގެ ނަންބަރު ދަށަންނަން ފެންނަނީ މާލެ އާއި ރާއްޖޭގެ ބަނިތަކުންނެވެ. ފަލިކާ ގެ ސަބަބުން ޕިންޓައިލް ގެ ނަންބަރު ދަށަންނަން ފެންނަނީ މާލެ އާއި ރާއްޖޭގެ ބަނިތަކުންނެވެ. ފަލިކާ ގެ ސަބަބުން ޕިންޓައިލް ގެ ނަންބަރު ދަށަންނަން ފެންނަނީ މާލެ އާއި ރާއްޖޭގެ ބަނިތަކުންނެވެ. ފަލިކާ ގެ ސަބަބުން ޕިންޓައިލް ގެ ނަންބަރު ދަށަންނަން ފެންނަނީ މާލެ އާއި ރާއްޖޭގެ ބަނިތަކުންނެވެ. ފަލިކާ ގެ ސަބަބުން ޕިންޓައިލް ގެ ނަންބަރު ދަށަންނަން ފެންނަނީ މާލެ އާއި ރާއްޖޭގެ ބަނިތަކުންނެވެ. ފަލިކާ ގެ ސަބަބުން ޕިންޓައިލް ގެ ނަންބަރު ދަށަންނަން ފެންނަނީ މާލެ އާއި ރާއްޖޭގެ ބަނިތަކުންނެވެ. ފަލިކާ ގެ ސަބަބުން ޕިންޓައިލް ގެ ނަންބަރު ދަށަންނަން ފެންނަނީ މާލެ އާއި ރާއްޖޭގެ ބަނިތަކުންނެވެ. ފަލިކާ ގެ ސަބަބުން ޕިންޓައިލް ގެ ނަންބަރު ދަށަންނަން ފެންނަނީ މާލެ އާއި ރާއްޖޭގެ ބަނިތަކުންނެވެ.



گھونٹا چھوڑنے والا

Common Ringed Plover

Charadrius hiaticula



توٹڈ ڈک

Tufted Duck

Aythya fuligula



عقاز قمریہ

Whiskered Tern

Chlidonias hybrida



تائید سوری

White-cheeked Tern

Sterna repressa



نارسه دوگالی

Roseate Tern

Sterna dougallii



سہیل

Common Tern

Sterna hirundo



تیر

Bridled Tern

Onychoprion anaethetus



مونتگُو ہاریر

Montagu's Harrier

Circus pygargus

ފަވަ ދަރަސާރު

Montagu's Harrier

Circus pygargus

Status: Northern winter visitor

Local Abundance: Rare

IUCN Red List Status: Least Concern (2021)

CITES Appendix: Not listed

Description: Measures between 39-49 cm in length. Males have balck bands across the wings and wingtips are black. Have narrow white patches on lower back. Males have mostly black plumage. Females are a rich chocolate color and have grey underwings. Juveniles are darker on top with reddish undersides.

މަގު ފަވަ ދަރަސާރު 39-49 ސ.މ. ގެ ދිގު ގަޅުގެ ތެރޭގައި ދެފަންނަނެވެ. ފަވަ ދަރަސާރުގެ ބުރަކަށި ވަށައިގެން ގަލަނި ލައިގެން ފެންނަނެވެ. ފަވަ ދަރަސާރުގެ ބުރަކަށި ވަށައިގެން ގަލަނި ލައިގެން ފެންނަނެވެ. ފަވަ ދަރަސާރުގެ ބުރަކަށި ވަށައިގެން ގަލަނި ލައިގެން ފެންނަނެވެ. ފަވަ ދަރަސާރުގެ ބުރަކަށި ވަށައިގެން ގަލަނި ލައިގެން ފެންނަނެވެ. ފަވަ ދަރަސާރުގެ ބުރަކަށި ވަށައިގެން ގަލަނި ލައިގެން ފެންނަނެވެ. ފަވަ ދަރަސާރުގެ ބުރަކަށި ވަށައިގެން ގަލަނި ލައިގެން ފެންނަނެވެ. ފަވަ ދަރަސާރުގެ ބުރަކަށި ވަށައިގެން ގަލަނި ލައިގެން ފެންނަނެވެ. ފަވަ ދަރަސާރުގެ ބުރަކަށި ވަށައިގެން ގަލަނި ލައިގެން ފެންނަނެވެ. ފަވަ ދަރަސާރުގެ ބުރަކަށި ވަށައިގެން ގަލަނި ލައިގެން ފެންނަނެވެ.



وَرْدِ قَه سَرِ سَر

Greater Sand Plover
Charadrius leschenaultii



قوچ پرنده

Ruff

Calidris pugnax



پرنده مهاجر

Eurasian Oystercatcher

Haematopus ostralegus



مَرَّوَجِيَّ عَسَلِيَّةٍ

European Bee-eater

Merops apiaster



درختان و پرندگان

European Roller

Coracias garrulus



۲۹۳

Great Frigatebird

Fregata minor



۳۳۳

Flesh-footed Shearwater

Ardenna carneipes



خبر رُکھ سرنے

Isabelline Wheatear

Oenanthe isabellina



جوانین پتrel

Jouanin's Petrel

Bulweria fallax

މިނި ފޮރިމް

Jouanin's Petrel

Bulweria fallax

- Status: Offshore migrant
- Local Abundance: Uncommon
- IUCN Red List Status: Near Threatened (2018)
- CITES Appendix: Not listed

Description: Typically measures between 30-32 cm in length. Sooty brown to blackish brown coloured plumage; tips of feathers near the bill may appear slightly lighter and greyer. Has distinctive wedge-shaped tail. A faint pale band may be apparent on the central upperwing of older birds. Males and females look alike, with juveniles resembling adults.

މިނި ފޮރިމް 32-30 ސެންޓިމީޓަރު ގެ ބުރުވަތައް ގެ ގޮތުގައި ހުރެއެވެ. މި ފޮރިމް ރަޖިންނު ފާލަންގެ ވަރަށް ފަސޭހައި ފެންނަން ހުރެއެވެ. މި ފޮރިމް ފިރިމީހާގެ ވަރަށް ފަސޭހައި ފެންނަން ހުރެއެވެ. މިނި ފޮރިމް ގެ ގޮތުގައި ހުރެއެވެ. މި ފޮރިމް ގެ ބުރުވަތައް ނިގޫ ގޮތުގައި ހުރެއެވެ. މިނި ފޮރިމް ގެ ބުރުވަތައް ނިގޫ ގޮތުގައި ހުރެއެވެ. މިނި ފޮރިމް ގެ ބުރުވަތައް ނިގޫ ގޮތުގައި ހުރެއެވެ. މިނި ފޮރިމް ގެ ބުރުވަތައް ނިގޫ ގޮތުގައި ހުރެއެވެ.



شیر تاج‌خروش

Brown Shrike

Lanius cristatus



کُسرِ سائمنوڈاکٹریر

Brown Skua

Stercorarius antarcticus



دروغہ سرخ گلوں کے درختوں پر

Red-Throated Pipit

Anthus cervinus



گُرُورِ سَوْتِی

Sooty Gull

Ichthyaetus hemprichii



قلمرو پرندگانی پاکستان

Chestnut-winged Cuckoo

Clamator coromandus



کھنڈی نڈری پر رگھو پر

Sykes's Short-toed Lark

Calandrella dukhunensis



بھنگی

Brown Noddy

Anous stolidus



پرنس پتیلورینس / پرنس پتیلورینس

Oriental Honey-Buzzard

Pernis ptilorhyncus



މާލެ ސަރަޙައްދުގެ ސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކޮށްފައިވާ ދިވެހިރާއްޖޭގެ ޕްރޮޓެކްޓެޑް އެރިއާތަކުގެ ތެރެއިން

Central Maldivian Heron

Butorides striata didii

ދަރިކަލުގެ ނަންބަރު 15 16 17 18 19 20
މަދަރުކަލުގެ ނަންބަރު 21 22 23 24 25 26

Central Maldivian Heron

Butorides striata didii

Status: Breeding resident

Local Abundance: Common

IUCN Red List Status: Unknown

CITES Appendix: Not listed

Description: Has a dark grey plumage with white underparts. Has a short neck with the chest and neck being white with a reddish-brown hue. Has distinctive bright yellow eyes and pale yellow legs. The beak is sharp and is relatively long.

ދަރިކަލު

މަދަރުކަލުގެ ނަންބަރު 15 16 17 18 19 20
މަދަރުކަލުގެ ނަންބަރު 21 22 23 24 25 26
މަދަރުކަލުގެ ނަންބަރު 27 28 29 30 31 32
މަދަރުކަލުގެ ނަންބަރު 33 34 35 36 37 38



جہازدار قمری

Caspian Tern

Hydroprogne caspia



نورسوز پرندہ

Green Sandpiper

Tringa ochropus



ہندوستان کی پرندوں کی فہرست

Indian Cormorant

Phalacrocorax fuscicollis



نُصْرَدُ الْفَرَسِ

Lesser Golden Plover

Pluvialis dominica



دندرونانٹھ انڈیکس

Forest Wagtail
Dendronanthus indicus



پوسکرینر

Common Sandpiper

Actitis hypoleucos



دریغ سہیلہ کے لیے

Wood Sandpiper

Tringa glareola



نوسنگ سونے چھریاں / ستر ستر

Common Hoopoe

Upupa epops



چوہدری

Sand Martin

Riparia riparia



پالید سلیٹ

Pallid Swift

Apus pallidus

ދިވެހިސަރުކާރުގެ ގެޒެޓްގައި ބަޔާންކޮށްފައިވާ ދިވެހިރާއްޖޭގެ ޖަނަވާތަކާއި ފަތިހުގެ ޖަނަވާތަކުގެ ނަންބަރުތަކުގެ ތެރޭގައި ހިމެނޭ ޖަނަވާތަކުގެ ނަންބަރުތަކެވެ.

Pallid Swift

Apus pallidus

Status: -

Local Abundance: Vagrant

IUCN Red List Status: Least Concern (2018)

CITES Appendix: Not Listed

Description: Large Swift measuring about 16 cm in length. Has broad, blunt wing tips and a short, shallowly forked tail. Has a broad head and robust body compared to other swifts. Overall brown in colour with a distinctive pale, triangular throat patch.

ދިވެހިސަރުކާރުގެ ގެޒެޓްގައި ބަޔާންކޮށްފައިވާ ދިވެހިރާއްޖޭގެ ޖަނަވާތަކާއި ފަތިހުގެ ޖަނަވާތަކުގެ ނަންބަރުތަކުގެ ތެރޭގައި ހިމެނޭ ޖަނަވާތަކުގެ ނަންބަރުތަކެވެ.

ދިވެހިސަރުކާރުގެ ގެޒެޓްގައި ބަޔާންކޮށްފައިވާ ދިވެހިރާއްޖޭގެ ޖަނަވާތަކާއި ފަތިހުގެ ޖަނަވާތަކުގެ ނަންބަރުތަކުގެ ތެރޭގައި ހިމެނޭ ޖަނަވާތަކުގެ ނަންބަރުތަކެވެ. ދިވެހިސަރުކާރުގެ ގެޒެޓްގައި ބަޔާންކޮށްފައިވާ ދިވެހިރާއްޖޭގެ ޖަނަވާތަކާއި ފަތިހުގެ ޖަނަވާތަކުގެ ނަންބަރުތަކުގެ ތެރޭގައި ހިމެނޭ ޖަނަވާތަކުގެ ނަންބަރުތަކެވެ.



سائبر

Common Swift

Apus apus



ہرنداپس کڈاکٹس

White-throated Needletail

Hirundapus caudacutus



ہندوستانی چوہدری

Indian Swiftlet
Aerodramus unicolor



سُرُفَره پَاسِفِیْک

Pacific Swift

Apus pacificus



پتروچیلڈن

Streak-throated Swallow

Petrochelidon fluvicola



مردو سرتر چویرنا

Red-rumped Swallow

Cecropis daurica

ދިވެހިސަރުކާރުގެ ގެޒެޓް

Red-rumped Swallow

Cecropis daurica

Status: Northern winter visitor

Local Abundance: Rare

IUCN Red List Status: Least Concern (2024)

CITES Appendix: Not Listed

Description: Medium sized bird measuring 16-17 cm in length. Has a robust build and flies with a slow, graceful motion. Has a long, deeply forked tail, with glossy blue-black feathers on its upper body, contrasting against a rich dark rufous rump. The underparts are white with bold black streaks, particularly on the throat. Juveniles look similar to adults but are generally browner, duller, and have a paler rump.

މަލްދިވެހިސަރުކާރުގެ ގެޒެޓް

މަލްދިވެހިސަރުކާރުގެ ގެޒެޓް 17-16 ސަސްޓިޓުޓްޔާއި ގުޅިގެން
 ސަރުކާރުގެ ޖަލްދުގައި ލިޔެފައިވާ ގޮތުން ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި
 ބަޔާންކޮށްފައިވާ ގޮތުން ފޮޓޯޓެޔް ކުރެއްވި ފޮޓޯޓެޔް ސަރުކާރުގެ
 ފޮޓޯޓެޔް ގޮތުން ފޮޓޯޓެޔް ކުރެއްވި ފޮޓޯޓެޔް ސަރުކާރުގެ ފޮޓޯޓެޔް
 ގޮތުން ފޮޓޯޓެޔް ކުރެއްވި ފޮޓޯޓެޔް ސަރުކާރުގެ ފޮޓޯޓެޔް ގޮތުން
 ފޮޓޯޓެޔް ކުރެއްވި ފޮޓޯޓެޔް ސަރުކާރުގެ ފޮޓޯޓެޔް ގޮތުން
 ފޮޓޯޓެޔް ކުރެއްވި ފޮޓޯޓެޔް ސަރުކާރުގެ ފޮޓޯޓެޔް ގޮތުން
 ފޮޓޯޓެޔް ކުރެއްވި ފޮޓޯޓެޔް ސަރުކާރުގެ ފޮޓޯޓެޔް ގޮތުން



گری تھروٹڈ مارٹن

Grey-throated Martin

Riparia chinensis



توتنگر مارش

Marsh Sandpiper

Tringa stagnatilis



تَیْرِ سِیِّ

Persian Shearwater

Puffinus persicus



پرنده‌های مهاجر در ایران

Black-winged Kite

Elanus caeruleus



چرزین سائون

Common Starling

Sturnus vulgaris



پرنده مهاجر

Pintail Snipe

Gallinago stenura



سُوْنُو اَتَمُو سَرُو مَرُو تَرُو

Swinhoe's Snipe

Gallinago megala



پتیرے اور پرندوں کی دیگر اہم ترین خصوصیات

Yellow Bittern

Ixobrychus sinensis



پرنده مهاجر

Tree Pipit

Anthus trivialis



چتر سبزه‌رنگ

Amur Falcon

Falco amurensis



مَرَسَمَةُ دُرَّاسِ

Purple Heron

Ardea purpurea



پتھر پتھر کے پتھر پتھر کے پتھر پتھر کے پتھر

Wilson's Storm-petrel

Oceanites oceanicus



پتھر پتھر کی طرح / سب سے پہلے اور سب سے آخر

White Tern

Gygis alba



سكوا جنوبى

South Polar Skua

Stercorarius maccormicki



މަލްދީވިއްޔާގެ ލިޔަން ހެރަން

Maldivian Little Heron

Butorides striata albidula



ދިވެހިސަރުކާރުގެ ގެޒެޓް ގައި ބަޔާންކުރި ދިވެހިގެޒެޓް

Maldivian Water Hen

Amaurornis phoenicurus maldivus


ދިވެހި ނާރުގެ ޗެލި

Maldivian Water Hen

Amaurornis phoenicurus maldivus

Status:	Breeding resident
Local Abundance:	Common
IUCN Red List Status:	Not Assessed
CITES Appendix:	Not listed
Description:	Small bird with bluish-black upperparts with a white face, throat and chest. Underparts has a reddish-brown tinge. The bill is of a greenish-yellow colour. Has yellow legs with relatively long toes.

ދިވެހި ނާރުގެ ޗެލި ޖަހާ ލަވަ ވަނީ ހަތަރު ފަދަ ޕެޝަކަތަކެވެ. ފަނޑު ނާރުގެ ޗެލި ޖަހާ ލަވަ ވަނީ ހަތަރު ފަދަ ޕެޝަކަތަކެވެ. ފަނޑު ނާރުގެ ޗެލި ޖަހާ ލަވަ ވަނީ ހަތަރު ފަދަ ޕެޝަކަތަކެވެ. ފަނޑު ނާރުގެ ޗެލި ޖަހާ ލަވަ ވަނީ ހަތަރު ފަދަ ޕެޝަކަތަކެވެ.



میریٹھ کے پرندوں کی فہرست

Oriental Pratincole

Glareola maldivarum



پرفیو ریسرچ

Audubon's Shearwater

Puffinus lherminieri

ދިވެހިރާއްޖޭގެ ޖަނަވާތަކާ ބެހޭ އިދާރާތަކުން ބަލާފައިވާ ރިޔާސީ ނަންބަރު: 12/2019/އ.ޖ.ޒ

Audubon's Shearwater

Puffinus lherminieri

Status:	Breeding resident
Local Abundance:	Common
IUCN Red List Status:	Least Concern (2018)
CITES Appendix:	Not listed

Description: Small bird measuring 27-36 cm in length. Has a mix of dark and white coloring. Wings are short and broad. Tail is rounded and long. Has a slim body and a large bill. Compared to similar species, is more white around the face and eyes and more dark feathers under the tail. Males and females appear similar but females are generally smaller. Some birds may have a brownish tint on the back.

ދިވެހިރާއްޖޭގެ ޖަނަވާތަކާ ބެހޭ އިދާރާތަކުން ބަލާފައިވާ ރިޔާސީ ނަންބަރު: 12/2019/އ.ޖ.ޒ



قزوینی قزوین

Eurasian Spoonbill

Platalea leucorodia



پرنیوں کی دنیا

Oriental Darter
Anhinga melanogaster



دروغہ سے بچنے کے لیے سچے سچے جانوروں کی کتاب

Long-tailed Shrike

Lanius schach



پرندۀ سرخ‌رُخ دم‌نارنگ

Long-tailed Skua
Stercorarius longicaudus

ކަރީ ސަރީ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ

Long-tailed Skua

Stercorarius longicaudus

Status: Offshore

Local Abundance: Rare

IUCN Red List Status: Least concern (2018)

CITES Appendix: Not listed

Description: Measures about 48-53 cm in length. In breeding season, adults plumage has greyish upperparts, black wings and a black tail. Has a white head with a brownish black cap and a yellowish neck. Central tail feathers are long and pointed. In non-breeding season, adults appear browner and do not have long tail feathers. Males and females look alike but males are more white on the underside than females. Juveniles are brown. Has light tips on the feathers and patterned brown and white bars on underwings and sides.

ބަލާ

ކަރީ ސަރީ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ 53-48
 ސަންދަންދެން ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ
 ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ
 ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ
 ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ
 ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ
 ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ
 ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ
 ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ
 ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ
 ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ ލަވަ



درومہ راجہ ♂ درومہ راجہ ♀

Crab-plover

Dromas ardeola



ہیمنٹوپس

Black-Winged Stilt
Himantopus himantopus

މަލްދީވުގެ ރާއްޖޭގެ ބައެއް ސަރަޙައްދުތަކުގައި ފެނިވާ ދިވެހި ޖަނަވާތަކުގެ ތެރެއިން ޖަނަވާތަކެކެވެ.

Black-Winged Stilt

Himantopus himantopus

- Status:** -
- Local Abundance:** Vagrant
- IUCN Red List Status:** Least Concern (2019)
- CITES Appendix:** Not listed

Description: Typically measures between 33-36 cm in length, with distinctively long pink legs and a long, straight black bill. Males have black upperparts with a greenish gloss and black wings with white underparts. Upperparts of females are dull brown and does not have a greenish gloss. Adults have black colouring on the back of the head and neck. Juveniles are similar to adult females .

މަލްދީވުގެ ރާއްޖޭގެ ބައެއް ސަރަޙައްދުތަކުގައި ފެނިވާ ދިވެހި ޖަނަވާތަކުގެ ތެރެއިން ޖަނަވާތަކެކެވެ. 36-33 ސެންޓިމީޓަރުގެ ދިގުކަން ހުންނަ ދިވެހި ޖަނަވާތަކެކެވެ. ފެނިވާ ދިވެހި ޖަނަވާތަކެކެވެ. ފެނިވާ ދިވެހި ޖަނަވާތަކެކެވެ. ފެނިވާ ދިވެހި ޖަނަވާތަކެކެވެ. ފެނިވާ ދިވެހި ޖަނަވާތަކެކެވެ. ފެނިވާ ދިވެހި ޖަނަވާތަކެކެވެ. ފެނިވާ ދިވެހި ޖަނަވާތަކެކެވެ. ފެނިވާ ދިވެހި ޖަނަވާތަކެކެވެ. ފެނިވާ ދިވެހި ޖަނަވާތަކެކެވެ. ފެނިވާ ދިވެހި ޖަނަވާތަކެކެވެ.



بڑا سفید چنگا

White-eyed Gull

Ichthyaetus leucophthalmus



پاپاسولا اباوٹی

Abbott's Booby

Papasula abbotti

ᐅᐃᐃᐃᐃᐃ
 𑊛𑊛𑊛𑊛𑊛

Abbott's Booby

Papasula abbotti

- Status: -
- Local Abundance: Vagrant
- IUCN Red List Status: Endangered (2019)
- CITES Appendix: Appendix I

Description: Measures about 79 cm in length. Has a thick serrated bill. Eyes are big. Head and neck are white. Has long feathers at the back which give it a shaggy appearance. Upper body is mostly white with black spots on the back and tail. Underside is white with black patches near the thighs. Bill is pale blue with a dark tip and legs are greyish-blue. Males and females look alike but females have a pinkish bill. Juveniles appear similar to adult males but have darker eyes.

𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 79 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛
 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛
 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛
 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛
 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛
 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛
 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛 𑊛𑊛𑊛



مَرَدَاقُ

Great Egret

Ardea alba



نم‌نورک‌ساز

Intermediate Egret

Ardea intermedia



دریغ سہولت

Steppe Gull

Larus fuscus barabensis



سبز سرخ گولڈ

Brown-headed Gull

Chroicocephalus brunnicephalus



گولک

Heuglin's Gull


Larus fuscus heuglini



۴۲۳۳ ۳۱۶

Black-Headed Gull

Chroicocephalus brunnicephalus



دروغہ سے بچنے کے لیے

Barn Swallow

Hirundo rustica



فہرست پرندوں کے نام / سائنس دانوں کے نام

Greater Flamingo

Phoenicopterus roseus



سنگھڑی

Common Kestrel


Falco tinnunculus



سرخس سرخس

Oriental Hobby

Falco severus



سڈوئے ٹائٹوڈ کٹرپر

Pomarine Skua

Stercorarius pomarinus



سڈسے بڈر

Northern Shoveler

Spatula clypeata



سائبر ڈیڈ ڈیڈ

Common Cuckoo

Cuculus canorus



سومتری کتے

Brahminy Kite

Haliastur indus



سائنس دانوں کے لیے

Common Coot

Fulica atra

Common Coot

Common Coot

Fulica atra

Status: -

Local Abundance: Vagrant

IUCN Red List Status: Least concern (2019)

CITES Appendix: Not listed

Description: Medium-sized waterbird about 36-38 cm in length. Has a plump, dark slate-grey to black body with a characteristic white frontal shield and bill that stands out against a dark head. Legs are greenish-grey with large lobed toes adapted for swimming. Shows a white trailing edge on the wings when in flight. Males and females look alike. Juveniles are duller grey with a smaller, pale bill and lack the prominent white frontal shield of adults.

Common Coot

Common Coot (Fulica atra) is a medium-sized waterbird, approximately 36-38 cm in length. It has a dark slate-grey to black body with a characteristic white frontal shield and a white bill that stands out against the dark head. The legs are greenish-grey and have large lobed toes adapted for swimming. In flight, it shows a white trailing edge on the wings. Males and females look alike. Juveniles are a duller grey with a smaller, pale bill and lack the prominent white frontal shield of adults.



پرنیوں کی دنیا

Western Spotted Dove

Spilopelia suratensis



اُرتولان بونینگ

Ortolan Bunting

Emberiza hortulana



پرنده مهاجر

Rosy Starling

Pastor roseus



جنگل کی پرند

Jungle Nightjar
Caprimulgus indicus



دکو بھنگو

Lesser Whistling Duck

Dendrocygna javanica



پنجاب کے پرندوں کی فہرست

Common Greenshank

Tringa nebularia

Common Greenshank

Common Greenshank

Tringa nebularia

Status:	Northern winter visitor
Local Abundance:	Common
IUCN Red List Status:	Least Concern (2016)
CITES Appendix:	Not Listed

Description: Typically measures between 30-35 cm in length. Characterised by a long, sturdy bill that tilts slightly upwards and long dull green legs. During breeding season, upperparts of the adults are streaked with black-brown and the head, neck and upper chest have darker streaks. In non-breeding season, adults have uniform grey upperparts, with a white face, neck and chest. Juveniles look similar to non-breeding adults but are more brown on top with buff edges on the feathers and a more streaked neck and chest.

Common Greenshank (Dhivehi: ޤިލިޞް ޖްރީންޝަންކް) is a species of bird in the family Scolopacidae. It is a northern winter visitor to the Maldives. The bird is typically 30-35 cm in length. It has a long, sturdy bill that tilts slightly upwards and long, dull green legs. During the breeding season, the upperparts of the adults are streaked with black-brown, and the head, neck, and upper chest have darker streaks. In the non-breeding season, the adults have uniform grey upperparts, a white face, neck, and chest. Juveniles look similar to non-breeding adults but are more brown on top with buff edges on the feathers and a more streaked neck and chest.



دردی سہ ماہی کے لیے سب سے زیادہ مناسب اور آسان کتاب

Spotted Redshank

Tringa erythropus



پرنیوں کی فہرست

Common Rosefinch
Carpodacus erythrinus

Index

ދިވެހި ދަނޑުގެ ވިޔަފާރި

CITES Appendix	IUCN Red List Status	Local Abundance	Status	Scientific Name	Common Name	ދިވެހި ނަންމު	
Not listed	Least Concern (2018)	Vagrant	-	Motacilla alba (Linnaeus, 1758)	White Wagtail	ހުދު ބަނޑުގެ ވިޔަފާރި	18
Not listed	Least Concern (2024)	Rare	Northern winter visitor	Terpsiphone paradisi (Linnaeus, 1758)	Indian Paradise-flycatcher	ހިންދު ދަނޑުގެ ވިޔަފާރި	20
Not listed	Least Concern (2020)	Rare	Northern winter visitor	Oriolus kundoo (Sykes, 1832)	Indian Golden Oriole	ހިންދު ދަނޑުގެ ވިޔަފާރި	22
Not listed	Least Concern (2018)	Vagrant	-	Saxicoloides fulicatus (Linnaeus, 1766)	Indian Robin	ހިންދު ދަނޑުގެ ވިޔަފާރި	24
Not listed	Least Concern (2020)	Vagrant	-	Coracina macei (Lesson, 1831)	Indian Cuckooshrike	ހިންދު ދަނޑުގެ ވިޔަފާރި	26
Not listed	Least concern (2024)	Vagrant	-	Coracias benghalensis (Linnaeus, 1758)	Indian Roller	ހިންދު ދަނޑުގެ ވިޔަފާރި	28
Not Listed	Least Concern (2018)	Vagrant	-	Cuculus micropterus (Gould, 1838)	Indian Cuckoo	ހިންދު ދަނޑުގެ ވިޔަފާރި	30
Not listed	Least Concern (2018)	Common	Non-breeding visitor	Fregata ariel (Grey, 1845)	Lesser Frigatebird	ހިންދު ދަނޑުގެ ވިޔަފާރި	32
Not listed	Unknown	Common	Breeding resident in south	Ardeola greyii phillipsi (Scheer, 1960)	Maldivian Pond Heron	ހިންދު ދަނޑުގެ ވިޔަފާރި	34
Appendix II	Near Threatened (2021)	Rare	North Winter Visitor	Circus macrourus (Gmelin, 1770)	Pallid Harrier	ހިންދު ދަނޑުގެ ވިޔަފާރި	36
Not listed	Least Concern (2020)	Rare	Straggler	Pelecanus onocrotalus (Linnaeus, 175)	Great White Pelican	ހިންދު ދަނޑުގެ ވިޔަފާރި	38
Not listed	Least Concern (2019)	Common	Breeding resident	Phaethon lepturus (Daudin, 1802)	White-Tailed Tropicbird	ހިންދު ދަނޑުގެ ވިޔަފާރި	40
Not listed	Least Concern (2018)	Vagrant	-	Hydrobates castro (Harcourt, 1851)	Band-rumped Storm-petrel	ހިންދު ދަނޑުގެ ވިޔަފާރި	42
Not Listed	Least Concern (2018)	Rare	Offshore migrant, Southern Winter	Pelagodroma marina (Latham, 1790)	White-faced Storm-petrel	ހިންދު ދަނޑުގެ ވިޔަފާރި	44
Not listed	Least Concern (2018)	Uncommon	Straggler	Sula dactylatra (Lesson, 1831)	Masked Booby	ހިންދު ދަނޑުގެ ވިޔަފާރި	46
Not listed	Vulnerable (2018)	Vagrant	-	Hydrobates leucorhous (Vieillot, 1818)	Leach's Storm-petrel	ހިންދު ދަނޑުގެ ވިޔަފާރި	48

Not listed	Near Threatened (2018)	Rare	Offshore Migrant	Calonectris leucomelas (Temminck, 1836)	Streaked Shearwater	މަލިކު ފަލުކު	50
Not listed	Near Threatened (2019)	Rare	Offshore migrant, April	Ardenna grisea (Gmelin, 1789)	Sooty Shearwater	ފަލުކު ފަލުކު	52
Not listed	Least Concern (2024)	Unknown	-	Ardea melanocephala (Children & Vigors, 1826)	Black-headed Heron	ކަލަ ފަލުކު	54
Not listed	Least concern (2024)	Rare	Northern winter visitor	Merops philippinus (Linnaeus, 1767)	Blue-tailed Bee-eater	ފަލުކު ފަލުކު ފަލުކު	56
Not listed	Least concern (2024)	Rare	Irruptive	Merops persicus (Pallas, 1773)	Blue-cheeked Bee-eater	ފަލުކު ފަލުކު ފަލުކު	58
Not Listed	Vulnerable (2019)	Vagrant	-	Streptopelia turtur (Linnaeus, 1758)	European Turtle Dove	ކަލަ ފަލުކު	60
Not listed	Least Concern (2024)	Uncommon	Northern winter visitor	Delichon urbicum (Linnaeus, 1758)	Northern House Martin	ކަލަ ފަލުކު	62
Not listed	Least Concern (2016)	Vagrant	-	Botaurus stellaris (Linnaeus, 1758)	Great Bittern	ކަލަ ފަލުކު	64
Not listed	Least Concern (2016)	Unknown	-	Dendrocygna bicolor (Vieillot, 1816)	Fulvous Whistling Duck	ކަލަ ފަލުކު	66
Not listed	Least Concern (2018)	Uncommon	Northern winter passage migrant	Motacilla flava (Linnaeus, 1758)	Western Yellow Wagtail	ކަލަ ފަލުކު	68
Not listed	Least Concern (2019)	Vagrant	-	Motacilla citreola (Pallas, 1776)	Citrine Wagtail	ކަލަ ފަލުކު	70
Appendix II	Least concern (2021)	Vagrant	-	Otus brucei (Hume, 1872)	Pallid Scops Owl	ކަލަ ފަލުކު	72
Not listed	Least Concern (2016)	Uncommon	May breed in south	Nycticorax nycticorax (Linnaeus, 1758)	Black-crowned Night Heron	ކަލަ ފަލުކު	74
Not listed	Least Concern (2024)	Regular	Northern winter visitor	Ardeola greyii greyii	Indian Pond Heron	ކަލަ ފަލުކު	76
Appendix II	Least Concern (2021)	Rare	Northern winter visitor	Falco naumanni (Fleischer, 1818)	Lesser Kestrel	ކަލަ ފަލުކު	78
Not listed	Least Concern (2024)	Vagrant	-	Lanius phoenicuroides (Schalow, 1875)	Red-tailed Shrike	ކަލަ ފަލުކު	80
Not Listed	Least Concern (2020)	Rare	Straggler	Phaethon rubricauda (Boddaert, 1783)	Red-tailed Tropicbird	ކަލަ ފަލުކު	82
Not listed	Least concern (2024)	Rare	Straggler	Botaurus cinnamomeus (Gmelin, 1789)	Cinnamon Bittern	ކަލަ ފަލުކު / ފަލުކު ފަލުކު	84
Not listed	Near Threatened (2019)	Rare	Northern winter visitor	Aythya nyroca (Güldenstädt, 1770)	Ferruginous Pochard	ކަލަ ފަލުކު	86
Not listed	Near Threatened (2024)	Common	Norther winter, some overstay	Arenaria interpres (Linnaeus, 1758)	Ruddy Turnstone	ކަލަ ފަލުކު	88
Not listed	Least Concern (2016)	Uncommon	Northern winter visitor	Tringa totanus (Linnaeus, 1758)	Common Redshank	ކަލަ ފަލުކު	90
Not listed	Least Concern (2020)	Uncommon	Non-breeding visitor	Sula sula (Linnaeus, 1766)	Red-footed Booby	ކަލަ ފަލުކު / ފަލުކު ފަލުކު	92

Not Listed	Least Concern (2019)	Rare	Straggler	Phaethon aethereus (Linnaeus, 1758)	Red-billed Tropicbird	سرخ بیل ٹروپک بڈ	94
Not listed	Least Concern (2018)	Common	Breeding resident	Sternula saundersi (Hume, 1877)	Saunders's Tern	سڈرس ٹرن	96
Not listed	Least concern (2019)	Uncommon	Northern winter visitor	Sternula albifrons (Pallas, 1764)	Little Tern	لیٹل ٹرن / سڈرس ٹرن	98
Not listed	Least Concern (2020)	Common	Offshore	Onychoprion fuscatus (Linnaeus, 1766)	Sooty Tern	سڈس ٹرن	100
Not listed	Least Concern (2016)	Uncommon	Northern winter visitor	Charadrius dubius (Scopoli, 1786)	Little Ringed Plover	لیٹل رینگڈ پلوور / سڈس ٹرن	102
Not listed	Least Concern (2018)	Vagrant	-	Charadrius asiaticus (Pallas, 1773)	Caspian Plover	کاسپین پلوور	104
Not listed	Vulnerable (2024)	Common	Northern winter visitor	Calidris ferruginea (Pontoppidan, 1763)	Curlew Sandpiper	کولرل سڈس ٹرن	106
Not listed	Least Concern (2016)	Rare	Northern winter visitor	Xenus cinereus (Güldenstädt, 1775)	Terek Sandpiper	تیرک سڈس ٹرن	108
Not listed	Least Concern (2024)	Common	North Winter Visitor	Numenius phaeopus (Linnaeus, 1758)	Whimbrel	ویمبریل	110
Not listed	Near Threatened (2016)	Uncommon	Northern winter visitor	Limosa lapponica (Linnaeus, 1758)	Bar-tailed Godwit	بار ٹیلڈ گڈ ویٹ	112
Not listed	Least concern (2024)	Vagrant	-	Threskiornis melanocephalus (Latham, 1790)	Black-headed Ibis	بلیک ہیڈڈ ایبیس	114
Not listed	Least Concern (2016)	Vagrant	-	Turdus obscurus (Gmelin, 1789)	Eyebrowed Thrush	ایبروڈڈ تھرسھ	116
Not listed	Least Concern (2016)	Uncommon	Northern winter visitor	Pluvialis fulva (Gmelin, 1789)	Pacific Golden Plover	پیسفک گولڈن پلوور	118
Not listed	Near Threatened (2019)	Vagrant	-	Egretta gularis (Bosc, 1792)	Western Reef Egret	سڈس ٹرن	120
Not listed	Near Threatened (2017)	Uncommon	North Winter Visitor	Numenius arquata (Linnaeus, 1758)	Eurasian Curlew	یوریشین کولرل	122
Not listed	Least Concern (2018)	Vagrant	-	Ichthyaetus ichthyaetus (Pallas, 1773)	Pallas's Gull	پالاس گولڈن	124
Not listed	Least Concern (2018)	Vagrant	-	Larus cachinnans (Pallas, 1811)	Yellow-legged Gull/Caspian Gull	یلو لیگڈ گولڈن	126
Appendix I	Least Concern (2021)	Rare	Northern winter visitor	Falco peregrinus (Tunstall, 1771)	Peregrine Falcon	پیریگرین فیلکن	128
Not listed	Least Concern (2018)	Common	Offshore migrant	Ardenna pacifica (Gmelin, 1789)	Wedge-Tailed Shearwater	ویڈج ٹیلڈ شیئر واٹر	130
Not listed	Least Concern (2018)	Common	Breeding resident	Thalasseus bergii (Lichtenstein, 1823)	Greater Crested Tern	گریٹر کریسٹڈ ٹرن	132
Appendix II	Least Concern (2021)	Uncommon	Northern winter visitor	Buteo buteo (Linnaeus, 1758)	Common Buzzard	کامن بڈزڈ	134
Not listed	Least Concern (2016)	Regular	Northern winter visitor	Anas querquedula (Linnaeus, 1758)	Garganey	گارجینی	136
Not listed	Least Concern (2019)	Rare	No recent records	Gelochelidon nilotica (Gmelin, 1789)	Gull-Billed Tern	گولڈن بیلڈ ٹرن	138
Not listed	Least concern (2024)	Uncommon	Northern winter visitor	Calidris alba (Pallas, 1764)	Sanderling	سڈرس ٹرن	140

Not Listed	Least Concern (2016)	Uncommon	Northern winter visitor	Charadrius alexandrinus (Linnaeus, 1758)	Kentish Plover	ކަލްޓިޝް ޕްލޮވަރ	142
Not listed	Least concern (2018)	Rare	Northern winter visitor	Calidris temminckii (Leisler, 1812)	Temminck's Stint	ޓެމިންކްސް ސްޓިންޓް	144
Not listed	Least Concern (2018)	Uncommon	Northern winter visitor	Calidris minuta (Leisler, 1812)	Little Stint	ލިޓްލް ސްޓިންޓް	146
Not listed	Near Threatened (2024)	Rare	Northern winter visitor	Calidris alpina (Linnaeus, 1758)	Dunlin	ޕްރިންސިޕަލް ސްޓިންޓް	148
Not listed	Least concern (2024)	Uncommon	Northern winter visitor	Calidris subminuta (von Middendorff, 1853)	Long-toed Stint	ލޮންގް-ޓޯޔްޑް ސްޓިންޓް	150
Not listed	Least concern (2016)	Vagrant	-	Glareola pratincola (Linnaeus, 1766)	Collared Pratincole	ކޮލަރެޑް ޕްރަޓިންކޮލް	152
Not Listed	Least Concern (2024)	Rare	Northern winter visitor	Streptopelia orientalis (Latham, 1790)	Oriental Turtle Dove	އޯރިއަންޓަލް ޓޯޓްލް ޑޯވް	154
Not listed	Least Concern (2020)	Uncommon	Northern winter visitor	Anas crecca (Linnaeus, 1758)	Common Teal	ކޮމަން ޓީލް	156
Not Listed	Least Concern (2018)	Common	Widespread	Anous tenuirostris (Temminck, 1823)	Lesser Noddy	ލެސަރ ނޮޑީ	158
Not listed	Least Concern (2018)	Vagrant	-	Ardena tenuirostris (Temminck, 1836)	Short-tailed Shearwater	ކޯޕްޕަރް ސްޕްރީޖް	160
Not listed	Least concern (2018)	Uncommon	Offshore, Northern Winter	Stercorarius parasiticus (Linnaeus, 1758)	Arctic Skua	އާރޯޕިކް ސްކޯޕް	162
Not listed	Least concern (2024)	Rare	Northern winter visitor	Oenanthe pleschanka (Lepechin, 1770)	Pied Wheatear	ޕީޑް ވްރީޓް ޕްލެޝްކާންކާ	164
Appendix II	Least concern (2021)	Rare	Irruptive	Asio flammeus (Pontoppidan, 1763)	Short-eared Owl	ކޯޕްޕަރް ޔަލްޕް ޕްލެއިންޓް	166
Not Listed	Least Concern (2016)	Introduced	Locally extinct?	Passer domesticus (Linnaeus, 1758)	House Sparrow	ހައުސް ސްޕަރޯ	168
Not listed	Least Concern (2018)	Common	Breeding Resident	Sterna sumatrana (Raffles, 1822)	Black-Naped Tern	ބަލަބް ޓަރުން	170
Not listed	Least concern (2016)	Rare	Northern winter visitor	Chlidonias leucopterus (Temminck, 1815)	White-winged Tern	ކްލައިޑޮނިޔާ ލީޔުކޮޕްޓަރޯ	172
Not listed	Least Concern (2018)	Uncommon	Offshore migrant, Northern winter	Bulweria bulwerii (Jardine & Selby, 1828)	Bulwer's Petrel	ބުލްވަރީ ޕެޓްރެލް	174
Not listed	Least Concern (2019)	Rare	Northern winter visitor	Plegadis falcinellus (Linnaeus, 1766)	Glossy Ibis	ގްލޯސީ އިބިސް	176
Appendix II	Least concern (2021)	Vagrant	-	Milvus migrans (Boddaert, 1783)	Black Kite	ބަލަބް ކައިޓް	178
Not listed	Near Threatened (2018)	Uncommon	Offshore migrant	Hydrobates monorhis (Swinhoe, 1867)	Swinhoe's Storm-petrel	ސުވިންހޯ ސްޓޯމް-ޕެޓްރެލް	180
Not listed	Least Concern (2024)	Rare	Northern winter visitor	Ixobrychus flavicollis (Latham, 1790)	Black Bittern	ބަލަބް ބިޓަރުން	182
Not listed	Least Concern (2016)	Uncommon	Northern winter visitor, may breed	Gallicrex cinerea (Gmelin, 1789)	Watercock	ވާޓަރކޮކް	184
Not listed	Least Concern (2018)	Uncommon	Straggler	Sula leucogaster (Boddaert, 1783)	Brown Booby	ބްރޯން ބޯބީ	186

Not listed	Least concern (2019)	Vagrant	-	Thalasseus sandvicensis (Latham, 1787)	Sandwich Tern	سانڈویچ ٹرن	188
Not listed	Least Concern (2016)	Common	Breeding Resident	Eudynamis scolopacea (Linnaeus, 1758)	Asian Koel	کھوکھو	190
Not Listed	Least Concern (2024)	Uncommon	Northern winter visitor	Nettapus coromandelianus (J.F. Gmelin, 1789)	Cotton Teal	سائبرینڈ	192
Not listed	Least Concern (2024)	Rare	Northern winter visitor	Alcedo atthis (Linnaeus, 1758)	Common Kingfisher	کومون کنگفشر	194
Not listed	Least Concern (2016)	Rare	Northern winter visitor	Apus affinis (J.E. Grey, 1830)	Little Swift	لیٹل سٹیفٹ	196
Not listed	Least Concern (2016)	Uncommon	Northern winter visitor	Egretta garzetta (Linnaeus, 1766)	Little Egret	لیٹل ایگریٹ	198
Appendix II	Least Concern (2021)	Rare	Northern winter visitor	Falco Subbuteo (Linnaeus, 1758)	Eurasian Hobby	لیٹل سببوتو	200
Not listed	Endangered (2023)	Uncommon	Northern winter visitor	Charadrius mongolus (Pallas, 1776)	Lesser Sand Plover	لیٹل ساند پلوور	202
Appendix II	Least Concern (2021)	Rare	North Winter Visitor	Pandion haliaetus (Linnaeus, 1758)	Osprey	اسپری	204
Appendix II	Least Concern (2021)	Uncommon	North Winter Visitor	Circus aeruginosus (Linnaeus, 1758)	Western Marsh Harrier	ایسٹرن مارش ہیریئر	206
Not listed	Near Threatened (2017)	Vagrant	-	Limosa limosa (Linnaeus, 1758)	Black-Tailed Godwit	بلیک ٹیلڈ گڈویٹ	208
Not listed	Least Concern (2016)	Unknown	-	Lymnocyptes minimus (Brunnich, 1764)	Jack Snipe	جیک سنیپ	210
Not listed	Least Concern (2016)	Uncommon	Northern winter visitor	Gallinago gallinago (Linnaeus, 1758)	Common Snipe	کومون سنیپ	212
Not listed	Least Concern (2019)	Vagrant	-	Vanellus gregarius (Pallas, 1771)	Sociable Lapwing	سوسیابل لاپونگ	214
Not listed	Least Concern (2019)	Common	Northern winter visitor	Bubulcus ibis (Linnaeus, 1758)	Cattle Egret	کیتل ایگریٹ	216
Not listed	Least Concern (2019)	Common	Breeding Resident	Ardea cinerea (Linnaeus, 1758)	Grey Heron	گری ہیرن	218
Not listed	Least Concern (2016)	Rare	Northern winter visitor	Motacilla cinerea (Tunstall, 1771)	Grey Wagtail	گری واگ ٹیل	220
Not listed	Near Threatened (2017)	Rare	Straggler	Pelecanus philippensis (Gmelin, 1789)	Spot-billed Pelican	اسپوٹ بیلڈ پیلکن	222
Not listed	Least Concern (2024)	Vagrant	-	Artamus fuscus (Vieillot, 1817)	Ashy Woodswallow	اشی وڈ سوا لو	224
Not Listed	Least Concern (2024)	Vagrant	-	Cacomantis passerinus (Vahl, 1797)	Grey-bellied Cuckoo	گری بیلڈ ککoo	226
Not listed	Least Concern (2019)	Uncommon	Breeds in far south	Gallinula chloropus (Linnaeus, 1758)	Common Moorhen	کومون مور ہین / ٹیل	228
Not listed	Least Concern (2018)	Common	Northern winter visitors, some overstay	Thalasseus bengalensis (Lesson, 1831)	Lesser Crested Tern	لیٹل کریسٹڈ ٹرن	230
Not listed	Least Concern (2018)	Vagrant	-	Oenanthe oenanthe (Linnaeus, 1758)	Northern Wheatear	نورن تھیٹیر	232
Not listed	Vulnerable (2024)	Uncommon	Northern winter visitor	Pluvialis Squatarola (Linnaeus, 1758)	Grey Plover	گری پلوور	234

Not listed	Least Concern (2019)	Uncommon	Northern winter visitor	Anas acuta (Linnaeus, 1758)	Northern Pintail	އިރިޔަލު ޖަލުބަނޑު	236
Not listed	Least Concern (2016)	Uncommon	Northern winter visitor	Charadrius hiaticula (Linnaeus, 1758)	Common Ringed Plover	އިރިޔަލު ޖަލުބަނޑު	238
Not listed	Least Concern (2016)	Uncommon	Northern winter visitor	Aythya fuligula (Linnaeus, 1758)	Tufted Duck	އިރިޔަލު ޖަލުބަނޑު	240
Not listed	Least Concern (2016)	Vagrant	-	Chlidonias hybrida (Pallas, 1811)	Whiskered Tern	ޅަލުބަނޑު	242
Not listed	Least concern (2019)	Vagrant	-	Sterna repressa (Hartert, 1916)	White-cheeked Tern	ޅަލުބަނޑު	244
Not listed	Least Concern (2018)	Uncommon	Breeding resident	Sterna dougallii (Montagu, 1813)	Roseate Tern	ޅަލުބަނޑު	246
Not listed	Least Concern (2018)	Common	Offshore Northern winter visitor	Sterna hirundo (Linnaeus, 1758)	Common Tern	ޅަލުބަނޑު	248
Not Listed	Least Concern (2019)	Common	Offshore	Onychoprion anaethetus (Scopoli, 1786)	Bridled Tern	ޅަލުބަނޑު	250
Appendix II	Least Concern (2021)	Rare	North Winter Visitor	Circus pygargus (Linnaeus, 1758)	Montagu's Harrier	ޅަލުބަނޑު	252
Not listed	Least Concern (2019)	Uncommon	Northern winter visitor	Charadrius leschenaultii (Lesson, 1826)	Greater Sand Plover	ޅަލުބަނޑު	254
Not listed	Least concern (2016)	Rare	Northern winter visitor	Calidris pugnax (Linnaeus, 1758)	Ruff	ޅަލުބަނޑު	256
Not listed	Near Threatened (2019)	Vagrant	-	Haematopus ostralegus (Linnaeus, 1758)	Eurasian Oystercatcher	ޅަލުބަނޑު	258
Not listed	Least Concern (2024)	Vagrant	-	Merops apiaster (Linnaeus, 1758)	European Bee-eater	ޅަލުބަނޑު	260
Not listed	Least Concern (2019)	Vagrant	-	Coracias garrulus (Linnaeus, 1758)	European Roller	ޅަލުބަނޑު	262
Not listed	Least Concern (2019)	Common	Non-breeding visitor	Fregata minor (Gmelin, 1789)	Great Frigatebird	ޅަލުބަނޑު	264
Not listed	Near Threatened (2018)	Common	South winter Offshore migrant	Ardenna carneipes (Gould, 1844)	Flesh-Footed Shearwater	ޅަލުބަނޑު	266
Not listed	Least Concern (2024)	Uncommon	Northern winter visitor	Oenanthe isabellina (Temminck, 1829)	Isabelline Wheatear	ޅަލުބަނޑު	268
Not listed	Near Threatened (2018)	Uncommon	Offshore migrant	Bulweria fallax (Jouanin, 1955)	Jouanin's Petrel	ޅަލުބަނޑު	270
Not listed	Least Concern (2024)	Rare	Northern winter visitor	Lanius cristatus (Linnaeus, 1758)	Brown Shrike	ޅަލުބަނޑު	272
Not listed	Least concern (2018)	Rare	Offshore, Southern Winter	Stercorarius antarcticus (Lesson, 1831)	Brown Skua	ޅަލުބަނޑު	274
Not listed	Least Concern (2018)	Rare	Northern winter visitor	Anthus cervinus (Pallas, 1811)	Red-Throated Pipit	ޅަލުބަނޑު	276
Not listed	Least concern (2018)	Vagrant	-	Ichthyætus hemprichii (Bruch, 1853)	Sooty Gull	ޅަލުބަނޑު	278
Not Listed	Least Concern (2024)	Vagrant	-	Clamator coromandus (Linnaeus, 1766)	Chestnut-winged Cuckoo	ޅަލުބަނޑު	280
Not listed	Least Concern (2024)	Rare	Northern winter visitor	Calandrella dukhunensis (Sykes, 1832)	Sykes's Short-toed Lark	ޅަލުބަނޑު	282

Not Listed	Least Concern (2020)	Common	Widespread	Anous stolidus (Linnaeus, 1758)	Brown Noddy	މުހަދަނި	284
Appendix II	Least Concern (2021)	Rare	Northern winter visitor	Pernis ptilorhynchus (Temminck, 1821)	Oriental Honey-Buzzard	ބަނޑު ބަނޑު / ބަނޑު ބަނޑު	286
Not listed	Unknown	Common	Breeding resident	Butorides striata didii (Phillips and Sims, 1958)	Central Maldivian Heron	މުހަދަނި ބަނޑު ބަނޑު	288
Not Listed	Least Concern (2018)	Rare	No recent records	Hydroprogne caspia (Pallas, 1770)	Caspian Tern	މުހަދަނި ބަނޑު ބަނޑު	290
Not listed	Least Concern (2016)	Vagrant	-	Tringa ochropus (Linnaeus, 1758)	Green Sandpiper	މުހަދަނި ބަނޑު ބަނޑު	292
Not listed	Least Concern (2018)	Vagrant	-	Phalacrocorax fuscicollis (Stephens, 1826)	Indian Cormorant	މުހަދަނި ބަނޑު ބަނޑު	294
Not listed	Least Concern (2024)	Unknown	-	Pluvialis dominica (Müller, 1776)	Lesser Golden Plover	މުހަދަނި ބަނޑު ބަނޑު	296
Not listed	Least Concern (2024)	Vagrant	-	Dendronanthus indicus (Gmelin, 1789)	Forest Wagtail	މުހަދަނި ބަނޑު ބަނޑު	298
Not listed	Least Concern (2016)	Common	Northern winter visitor	Actitis hypoleucos (Linnaeus, 1758)	Common Sandpiper	މުހަދަނި ބަނޑު ބަނޑު	300
Not listed	Least Concern (2016)	Uncommon	Northern winter visitor	Tringa glareola (Linnaeus, 1758)	Wood Sandpiper	މުހަދަނި ބަނޑު ބަނޑު	302
Not listed	Least concern (2020)	Rare	Northern winter visitor	Upupa epops (Linnaeus, 1758)	Common Hoopoe	މުހަދަނި ބަނޑު ބަނޑު / މުހަދަނި ބަނޑު ބަނޑު	304
Not listed	Least Concern (2016)	Uncommon	Autumn passage migrant	Riparia riparia (Linnaeus, 1758)	Sand Martin	މުހަދަނި ބަނޑު ބަނޑު	306
Not listed	Least Concern (2018)	Vagrant	-	Apus pallidus (Shelley, 1870)	Pallid Swift	މުހަދަނި ބަނޑު ބަނޑު	308
Not listed	Least Concern (2016)	Uncommon	Autumn passage(?)	Apus apus (Linnaeus, 1758)	Common Swift	މުހަދަނި ބަނޑު ބަނޑު	310
Not Listed	Least Concern (2024)	Rare	Northern winter visitor	Hirundapus caudacutus (Latham, 1801)	White-throated Needletail	މުހަދަނި ބަނޑު ބަނޑު	312
Not Listed	Least Concern (2024)	Rare	Northern winter visitor	Aerodramus unicolor (Jerdon, 1840)	Indian Swiftlet	މުހަދަނި ބަނޑު ބަނޑު	314
Not Listed	Least Concern (2024)	Rare	Northern winter visitor	Apus pacificus (Latham, 1801)	Pacific Swift	މުހަދަނި ބަނޑު ބަނޑު	316
Not listed	Least Concern (2024)	Vagrant	-	Petrochelidon fluvicola (Blyth, 1855)	Streak-throated Swallow	މުހަދަނި ބަނޑު ބަނޑު	318
Not listed	Least Concern (2024)	Rare	Northern winter visitor	Cecropis daurica (Laxmann, 1769)	Red-rumped Swallow	މުހަދަނި ބަނޑު ބަނޑު	320
Not listed	Least Concern (2024)	Vagrant	-	Riparia chinensis (Grey, 1830)	Grey-throated Martin	މުހަދަނި ބަނޑު ބަނޑު	322
Not listed	Least Concern (2016)	Uncommon	Northern winter visitor	Tringa stagnatilis (Bechstein, 1803)	Marsh Sandpiper	މުހަދަނި ބަނޑު ބަނޑު	324
Not listed	Least Concern (2018)	Vagrant	-	Puffinus persicus (Hume, 1872)	Persian Shearwater	މުހަދަނި ބަނޑު ބަނޑު	326
Appendix II	Least Concern (2019)	Rare	Breeding resident in south	Elanus caeruleus (Desfontaines, 1789)	Black-winged Kite	މުހަދަނި ބަނޑު ބަނޑު	328

Protected Birds of The Maldives

Not listed	Least Concern (2018)	Vagrant	-	<i>Sturnus vulgaris</i> (Linnaeus, 1758)	Common Starling	ދިވެހި ސަރުބަލި	330
Not listed	Least Concern (2016)	Uncommon	Northern winter visitor	<i>Gallinago stenura</i> (Bonaparte, 1831)	Pintail Snipe	އިތުރު ފަތިރި ދަނޑު ދަނޑު	332
Not listed	Least Concern (2016)	Vagrant	-	<i>Gallinago megala</i> (Swinhoe, 1861)	Swinhoe's Snipe	ފަތިރި ދަނޑު ދަނޑު	334
Not listed	Least Concern (2024)	Rare	Straggler	<i>Ixobrychus sinensis</i> (Gmelin, 1789)	Yellow Bittern	ދަނޑު ދަނޑު ސަރި	336
Not listed	Least Concern (2018)	Vagrant	-	<i>Anthus trivialis</i> (Linnaeus, 1758)	Tree Pipit	ދަނޑު ފަތިރި ދަނޑު	338
Appendix II	Least Concern (2021)	Uncommon	Autumn passage migrant	<i>Falco amurensis</i> (Radde, 1863)	Amur Falcon	ދަނޑު ސަރުބަލި	340
Not listed	Least concern (2019)	Rare	Northern winter visitor	<i>Ardea purpurea</i> (Linnaeus, 1766)	Purple Heron	ދަނޑު ދަނޑު	342
Not Listed	Least Concern (2018)	Common	Southern winter offshore migrant	<i>Oceanites oceanicus</i> (Kuhl, 1820)	Wilson's Storm-petrel	ދަނޑު ސަރުބަލި ސަރުބަލި ސަރުބަލި	344
Not listed	Least Concern (2022)	Common	Breeding resident in South	<i>Gygis alba</i> (Sparman, 1786)	White Tern	ދަނޑު ސަރުބަލި / ސަރުބަލި ދަނޑު	346
Not listed	Least concern (2018)	Rare	Offshore, Southern Winter	<i>Stercorarius maccormicki</i> (Saunders, 1893)	South Polar Skua	ދަނޑު ސަރުބަލި ސަރުބަލި ސަރުބަލި	348
Not listed	Unknown	Common	Breeding resident	<i>Butorides striata albidula</i> (Bangs, 1904)	Maldivian Little Heron	ދަނޑު ދަނޑު ސަރި	350
Not Listed	Not Assessed	Common	Breeding resident	<i>Amaurornis phoenicurus maldivus</i> (Pennant, 1769)	Maldivian Water Hen	ދަނޑު ސަރުބަލި	352
Not listed	Least concern (2024)	Uncommon	Northern winter visitor	<i>Glareola maldivarum</i> (Forster, 1795)	Oriental Pratincole	ދަނޑު ފަތިރި ދަނޑު	354
Not listed	Least Concern (2018)	Common	Breeding resident	<i>Puffinus lherminieri</i> (Lesson, 1839)	Audubon's Shearwater	ދަނޑު ސަރުބަލި	356
Appendix II	Least Concern (2016)	Rare	Northern winter visitor	<i>Platalea leucorodia</i> (Linnaeus, 1758)	Eurasian Spoonbill	ފަތިރި ދަނޑު ސަރި	358
Not listed	Least Concern (2024)	Vagrant	-	<i>Anhinga melanogaster</i> (Pennant, 1769)	Oriental Darter	ދަނޑު ފަތިރި ދަނޑު	360
Not listed	Least Concern (2024)	Vagrant	-	<i>Lanius schach</i> (Linnaeus, 1758)	Long-tailed Shrike	ދަނޑު ފަތިރި ދަނޑު ސަރި	362
Not listed	Least concern (2018)	Rare	Offshore	<i>Stercorarius longicaudus</i> (Vieillot, 1819)	Long-tailed Skua	ދަނޑު ސަރުބަލި ސަރުބަލި ސަރުބަލި	364
Not listed	Least Concern (2024)	Uncommon	Northern winter visitor	<i>Dromas ardeola</i> (Paykull, 1805)	Crab-plover	މަލިބަލި ފަތިރި ފަތިރި	366
Not listed	Least Concern (2019)	Vagrant	-	<i>Himantopus himantopus</i> (Linnaeus, 1758)	Black-Winged Stilt	ފަތިރި ފަތިރި ސަރި	368
Not listed	Least Concern (2018)	Vagrant	-	<i>Ichthyaeus leucophthalmus</i> (Temminck, 1825)	White-eyed Gull	ފަތިރި ފަތިރި ސަރި	440
Appendix I	Endangered (2019)	Vagrant	-	<i>Papasula abbotti</i> (Ridgway, 1893)	Abbott's Booby	ފަތިރި ފަތިރި ސަރި	370

Not listed	Least Concern (2016)	Uncommon	Northern winter visitor	Ardea alba (Linnaeus, 1758)	Great Egret	گرت ایگریٹ	374
Not listed	Least concern (2020)	Rare	Straggler	Ardea intermedia (Wagler, 1829)	Intermediate Egret	انٹرمیڈیٹ ایگریٹ	376
Not listed	Not listed	Rare	Northern winter visitor	Larus fuscus barabensis (Johansen, 1960)	Steppe Gull	سٹیپ گولڈ	378
Not listed	Least Concern (2018)	Vagrant	-	Chroicocephalus brunnicephalus (Jerdon, 1840)	Brown-Headed Gull	برائن ہیڈڈ گولڈ	380
Not listed	Least Concern (2018)	Rare	North Winter visitor	Larus fuscus heuglini (Bree, 1876)	Heuglin's Gull	ہیولین گولڈ	382
Not listed	Least Concern (2018)	Vagrant	-	Chroicocephalus brunnicephalus (Jerdon, 1840)	Black-Headed Gull	بلاک ہیڈڈ گولڈ	384
Not listed	Least Concern (2018)	Uncommon	Northern winter visitor	Hirundo rustica (Linnaeus, 1758)	Barn Swallow	بارن سوائلو	386
Appendix II	Least Concern (2019)	Rare	Northern winter visitor	Phoenicopterus roseus (Pallas, 1811)	Greater Flamingo	گریٹر فلیمینگو / سوائٹ ویٹر	388
Appendix II	Least Concern (2021)	Uncommon	Northern winter visitor	Falco tinnunculus (Linnaeus, 1758)	Common Kestrel	کسٹریل	390
Appendix II	Least Concern (2021)	Vagrant	-	Falco severus (Horsfield, 1821)	Oriental Hobby	اورینٹل ہابی	392
Not listed	Least concern (2018)	Uncommon	Offshore, Northern Winter	Stercorarius pomarinus (Temminck, 1815)	Pomarine Skua	پومارین سکیوا	394
Not listed	Least Concern (2019)	Regular	Northern winter visitor	Spatula clypeata (Linnaeus, 1758)	Northern Shoveler	نارٹرن شوولر	396
Not Listed	Least Concern (2024)	Uncommon	Autumn passage migrant(?)	Cuculus canorus (Linnaeus, 1758)	Common Cuckoo	کوکو	398
Appendix II	Least concern (2024)	Vagrant	-	Haliastur indus (Boddaert, 1783)	Brahminy Kite	براہمنی کیٹ	400
Not listed	Least concern (2019)	Vagrant	-	Fulica atra (Linnaeus, 1758)	Common Coot	کوکو	402
Not Listed	Least Concern (2024)	Vagrant	-	Spilopelia suratensis (Gmelin, 1789)	Western Spotted Dove	سپیلوپلیا سوراتینسیس	404
Appendix III	Least Concern (2016)	Vagrant	-	Emberiza hortulana (Linnaeus, 1758)	Ortolan Bunting	اورٹولان ہنٹنگ	406
Not listed	Least Concern (2016)	Rare	Northern winter visitor	Pastor roseus (Linnaeus, 1758)	Rosy Starling	روسے اسٹارلنگ	408
Not Listed	Least Concern (2024)	Vagrant	-	Caprimulgus indicus (Latham, 1790)	Jungle Nightjar	جنگل نائٹجر	410
Not Listed	Least Concern (2024)	Vagrant	-	Dendrocygna javanica (Horsfield, 1821)	Lesser Whistling Duck	لیسٹر ویسٹلنگ ڈک	412
Not listed	Least Concern (2016)	Common	Northern winter visitor	Tringa nebularia (Gunnerus, 1767)	Common Greenshank	گرن سھانک	414
Not listed	Least Concern (2016)	Vagrant	-	Tringa erythropus (Pallas, 1764)	Spotted Redshank	سپوٹڈ ریڈ سھانک	416
Appendix III	Least Concern (2024)	Vagrant	-	Carpodacus erythrinus (Pallas, 1770)	Common Rosefinch	ریس سھانک	418

ދިވެހިސަރުކާރުގެ ގެޒެޓް، ސަރުކާރުގެ ގާނޫނުތަކާއި ސަރުކާރުގެ ފަރާތުން ދެއްވާ ފަރާތްތަކުގެ ނަންބަރު

Ministry of Climate Change, Environment and Energy
Protected Birds of the Maldives



